


# Noisy surroundings take toll on short-term memory

September 6 2012



While participants kept the numbers they had heard in mind, alpha wave increased, peaking, when bad acoustics were combined with longer strings of numbers. Credit: Max Planck Society

Have you ever noticed how tiresome it can be to follow a conversation at a noisy party? Rest assured: this is not necessarily due to bad hearing – although that might make things worse. Scientists at the Max Planck Institute for Human Cognitive and Brain Sciences in Leipzig have found that adverse listening situations are difficult for the brain, partly because they draw on the same, limited resources supporting our short-term memory. The new findings are particularly relevant to understanding the cognitive consequences of hearing damage, a condition that affects an increasing number of people.

Whether we are engaged in small talk or trying to memorise a telephone number - it is our short-term memory that ensures we don't lose track.

But what if the very same memory gets additionally taxed because the words to be remembered are hard to understand? This is suggested by a new study conducted by Jonas Obleser and his team at the [Max Planck Research Group "Auditory Cognition"](#)

In the experiment, listeners were asked to memorise a few digits they heard (e.g., "2...5...9...3") for just over a second. This is a very easy task, and when asked whether they remembered hearing a certain digit, listeners answered correctly 90 percent of the time. During this process, Obleser and his colleagues were using magnetoencephalography to measure so-called "[alpha waves](#)" in the brain. "The brain tends to time its activity in rhythmic waves with the [alpha rhythm](#) consisting of 8 to 12 activity waves per second", Obleser explains. "The reason we were interested in this particular rhythm is that the strength of the alpha waves has been shown to indicate how much information you are currently storing in your short-term memory, with more vigorous alpha waves signalling a busier time."

As expected, alpha waves were stronger when [listeners](#) had more digits to memorise. But surprisingly, the strength of the alpha waves also depended on the acoustic format of the speech signal, which the scientists varied from clear to heavily degraded. The harder the digits were to understand, the stronger the alpha waves became. Speech content (e.g., a phone number you would like to remember) and speech acoustics (such as the noisy party at which you overheard that phone number) thus draw on a shared brain resource.

In our everyday lives our [short term memory](#) has a natural limit, which may be reached faster in noisy surroundings. These findings may be particularly relevant to populations who are constantly receiving such adverse input: People with hearing damage, or – most drastically – people who have restored but very fragmented hearing through an artificial inner ear, the so-called cochlear implant.

Further studies could examine in more detail how chronic sensory degradations affect the executive functions of the brain.

**More information:** Obleser, J., Woestmann, M., Hellbernd, N., Wilsch, A. , Maess, B. (2012). Adverse listening conditions and memory load drive a common alpha oscillatory network. *Journal of Neuroscience*. September 5, 2012, 32(36):12376 –12383

Provided by Max Planck Society

Citation: Noisy surroundings take toll on short-term memory (2012, September 6) retrieved 21 July 2023 from <https://medicalxpress.com/news/2012-09-noisy-toll-short-term-memory.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.