

## Sleep apnea common among stroke-related brainstem injuries

February 12 2014

People whose brainstems are affected by their stroke have a significantly higher prevalence of sleep apnea than those who have stroke-related injury elsewhere in the brain, according to research presented at the American Stroke Association's International Stroke Conference 2014.

Sleep apnea is marked by interrupted breathing during sleep and can lead to serious health problems including heart disease and <u>stroke</u>.

"This is the largest population-based study to address the issue of the location of the brain injury and its relationship to sleep apnea in poststroke <u>patients</u>," said Devin L. Brown, M.D., M.S., lead author of the study and associate professor of neurology and associate director of the stroke program at the University of Michigan in Ann Arbor.

The 355 <u>ischemic stroke</u> patients, average 65 years and 55 percent men, who participated in the study, were drawn from those already enrolled in the Brain Attack Surveillance in Corpus Christi (BASIC) Project. Hispanics represented 59 percent of the study group; Non-Hispanic whites 35 percent; Native Americans 1 percent; and African-Americans 4 percent.

According to researchers, patients were also offered sleep apnea screening with a portable respiratory monitor and were screened about 13 days after a stroke.

Neurologists interpreted CT and MRI brain imaging scan results to



determine if patients had brainstem involvement or no brainstem involvement.

Of the 11 percent of 355 stroke patients with brainstem injury, 84 percent had sleep apnea. Of those without brainstem involvement, 59 percent had sleep apnea.

"While these numbers are high, more research into the relationship between stroke and sleep apnea is needed before we recommend routine sleep apnea screening in post-<u>stroke patients</u>," Brown said.

In the future, Brown said her group will continue to explore the relationship between <u>sleep apnea</u> and stroke.

Provided by American Heart Association

Citation: Sleep apnea common among stroke-related brainstem injuries (2014, February 12) retrieved 20 December 2023 from <u>https://medicalxpress.com/news/2014-02-apnea-common-stroke-related-brainstem-injuries.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.