

Surgery improves survival rates for men with prostate cancer if radiation treatments fail

March 10 2016



Naveen Pokala, M.D., an assistant professor in the Division of Urology at the University of Missouri School of Medicine, performs a robotic surgery. Pokala found that a complex surgical procedure improves survival rates for men with prostate cancer after radiation therapy has failed. Credit: Justin Kelley/MU Health

Approximately 14 percent of men will be diagnosed with prostate cancer at some point in their lifetimes, according to the National Institutes of Health. Radiation therapy traditionally has been a primary treatment for the cancer, but one-fourth of men have a recurrence of prostate cancer within five years after the therapy. Now, a University of Missouri School of Medicine researcher has found that a complex procedure to remove the prostate achieves excellent long-term survival for men after radiation therapy has failed.

"Prostate cancer, unfortunately, is a common cancer, and more than 27,000 men are estimated to have died from the disease in 2015," said Naveen Pokala, M.D., an assistant professor in the Division of Urology at the MU School of Medicine and lead author of the study. "By studying a national database of prostate cancer cases, we found that a procedure known as salvage radical prostatectomy can greatly increase a man's chance of survival when traditional [radiation therapy](#) has failed to eradicate the cancer."

Using the Surveillance, Epidemiology and End Results (SEER) program database, Pokala and his research team studied 364 patients who underwent a salvage radical prostatectomy surgery after unsuccessful radiation treatments. Looking at survival rates, the researchers found that 88.6 percent of men were still alive 10 years later and 72.7 percent of men were still alive 20 years later.

During a salvage radical prostatectomy, the [prostate gland](#) and surrounding tissue are surgically removed to keep the cancer from spreading. The procedure is challenging because tissue that surrounds the prostate is scarred during radiation treatment, making it difficult for the surgeon to identify and cut out tissue that needs to be removed. If the cancer is localized, a highly skilled surgeon can remove the gland and surrounding tissue using a robotic minimally invasive technique or through an open surgery.

"Because radical prostatectomy is a complex surgery, there can be a reluctance to undergo the procedure," Pokala said. "However, this study shows that it is a viable treatment option. This can bring a renewed hope and peace of mind to [men](#) living with [prostate cancer](#)."

The study, "Survival Outcomes in Men Undergoing Radical Prostatectomy after Primary Radiation Treatment for Adenocarcinoma of the Prostate," recently was published by *Clinical Genitourinary Cancer*, a peer-reviewed journal on the detection, diagnosis, prevention and treatment of genitourinary cancers.

Provided by University of Missouri-Columbia

Citation: Surgery improves survival rates for men with prostate cancer if radiation treatments fail (2016, March 10) retrieved 24 November 2023 from <https://medicalxpress.com/news/2016-03-surgery-survival-men-prostate-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--