

Cancer mortality differs among Asian ethnic groups

October 3 2016

The growing Asian American population in the United States presents a diverse set of health behaviors and health outcomes, and may benefit from cancer prevention and screening efforts tailored to specific Asian ethnic groups, according to a study published in *Cancer Epidemiology, Biomarkers & Prevention*, a journal of the American Association for Cancer Research.

"Cancer is the second leading cause of death in the United States, yet trends and current patterns in the mortality burden of cancer among Asian American ethnic groups have not been documented," said the study's senior author, Latha P. Palaniappan, MD, MS, a clinical professor of medicine at Stanford University School of Medicine in California. "To fill this knowledge gap, we examined ethnicity-specific cancer mortality rates and trends during the years 2003-2011."

In recent years, Asian Americans have surpassed Hispanics as the most rapidly growing racial/ethnic group in the United States. There are currently 17.3 million Asian Americans in the U.S., representing 5.6 percent of the population. By the year 2050, the U.S. Census Bureau estimates the Asian American population will exceed 40 million.

Beginning in 2003, U.S. death certificates identified six major subgroups of the Asian population—Asian Indian, Chinese, Filipino, Japanese, Korean, and Vietnamese. Until the release of the new death certificate, only seven states collected detailed information on Asian ethnicity, which limited knowledge of how cancer affected each group, said

Caroline A. Thompson, PhD, MPH, an assistant professor of epidemiology in the Graduate School of Public Health at San Diego State University and the study's lead author.

For this study, Palaniappan, Thompson and colleagues obtained data on 85,616 cancer-related deaths from 2003 to 2011 from the National Center for Health Statistics. They examined mortality statistics by cancer type, sex, and ethnicity. Some key findings:

- Overall, Asian Americans experience lower cancer mortality rates than non-Hispanic whites, but certain exceptions point to differences in biological or lifestyle risk factors.
- Stomach cancer accounted for 10 to 15 percent of Korean cancer deaths, and 5 to 10 percent of Chinese, Japanese, and Vietnamese cancer deaths, compared with less than 2 percent of cancer deaths in non-Hispanic whites.
- Liver cancer accounted for 22 percent of cancer deaths in Vietnamese males, 12 percent in Chinese males, and 10 percent in Vietnamese females, compared with less than 2.5 percent of cancer deaths in non-Hispanic whites.
- Lung cancer is the leading cause of cancer-related death among men of all Asian American ethnicities, and for most women. However, the researchers noted that while lung cancer mortality trends are stable or declining among most of the population, there has been a modest increase in lung cancer deaths among Chinese, Filipina, and Japanese women.
- Breast cancer deaths accounted for 19.5 percent of all Filipina cancer deaths and 19.8 percent of Asian Indian cancer deaths, compared with 14.6 percent of cancer deaths among non-Hispanic white women. Among other Asian American women, breast cancer death rates were significantly lower than those for non-Hispanic white women.

"Our findings suggest that national [cancer prevention](#) efforts may benefit from ethnic-specific information on the cancers that affect each specific subgroup within the Asian American population," Thompson said. "For patients, it's important to understand your risk profile, and to consider cultural, lifestyle, or modifiable [health behaviors](#) that might increase your risk of cancer."

The authors suggested that screening efforts tailored to specific populations could be useful in preventing cancer or diagnosing it at an earlier stage. For example, they noted, gastric cancer screening programs are well established in Japan and South Korea, and could potentially benefit the Japanese and South Korean population living in America.

Such efforts would benefit significantly from the continued collection of data on specific Asian subgroups, they added.

"We believe this study highlights the importance of capturing and considering distinct Asian ethnicity in census data, disease surveillance and vital statistics data, as well as medical records and research studies," Palaniappan said.

Thompson said a limitation of the study is the potential for misclassification, since death certificates are generally completed by medical personnel who may not know detailed ethnicity.

More information: M. B. Gilkey et al. Quality of Physician Communication about Human Papillomavirus Vaccine: Findings from a National Survey, *Cancer Epidemiology Biomarkers & Prevention* (2015). [DOI: 10.1158/1055-9965.EPI-15-0326](https://doi.org/10.1158/1055-9965.EPI-15-0326)

Provided by American Association for Cancer Research

Citation: Cancer mortality differs among Asian ethnic groups (2016, October 3) retrieved 20 July 2023 from <https://medicalxpress.com/news/2016-10-cancer-mortality-differs-asian-ethnic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.