

Study duration affects rate of diabetes remission post-RYGB

October 29 2016



(HealthDay)—After Roux-en-Y gastric bypass (RYGB) surgery,



variations in rates of diabetes remission are primarily related to differences in the definition of remission and study duration, according to a review published online Oct. 13 in *Diabetes Care*.

Deanna J.M. Isaman, Ph.D., from the University of Michigan in Ann Arbor, and colleagues conducted a literature review that identified 10 large (n > 100), recent (index surgery since 2000) studies of diabetes remission after RYGB. Studies differed in definitions of remission, lengths of follow-up (one year versus at least three years), reported outcomes, and risks of attrition bias.

The researchers found that reported rates of partial remission were 10 to 30 percentage points higher than rates of complete remission. Sixty-nine percent of the variability in cumulative remission rates was explained by study duration. Variability up to 87 percent was explained by adjustment for attrition. Three-year cumulative, complete remission rates ranged from 63 to 65 percent, with adjustment for attrition. However, relapse was not accounted for in these results. Accounting for relapse, attritionadjusted, three-year prevalent complete remission rates were 23 percent.

"Future studies should report both cumulative and prevalent <u>remission</u> to aid decision making and more easily compare studies," the authors write.

More information: <u>Full Text (subscription or payment may be required)</u>

Copyright © 2016 HealthDay. All rights reserved.

Citation: Study duration affects rate of diabetes remission post-RYGB (2016, October 29) retrieved 11 July 2023 from https://medicalxpress.com/news/2016-10-duration-affects-diabetes-



remission-post-rygb.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.