


Sensory stimuli control dopamine in the brain

January 13 2017



A four-millimetre-long larvae of zebrafish with visible nerve cells under the microscope. Credit: Research group Driever

Regardless of whether we are sitting in a loud aeroplane or walking through a quiet forest clearing, how humans perceive their environment depends on the stimuli. This, in turn, affects our behaviour – sometimes consciously, sometimes subconsciously. In their study of fish larvae, Prof. Dr. Wolfgang Driever and his team of neurobiologists at the University of Freiburg have discovered that a group of nerve cells in the forebrain release the neurotransmitter dopamine when activated by tactile or certain visual stimuli. These dopaminergic nerve cells send connections to almost all parts of the brain and spinal cord, thereby affecting the functions of many circuits. These new findings could play a role in the future treatment of such illnesses as restless leg syndrome, a condition in which patients have unpleasant sensations in their limbs during sleep. The researchers have published their research results in the journal *Current Biology*.

For their research, the scientists studied the four-millimetre-long larvae of zebrafish, which are common aquarium fish. The scientists observed the activity of individual dopaminergic nerve cells within the brains of the larvae, which were alert and active, under a microscope. The researchers were able to make their activity visible using optogenetic calcium sensors, which emit light in active nerve cells. Until now, studies of the dopaminergic nerve cells in vertebrates have primarily focused on the midbrain, where the dopaminergic cells are involved in the control of locomotion and reward behaviour. These become functionally impaired in patients with Parkinson's disease. The dopaminergic neurons of the forebrain, on the other hand, have been little researched until now because they are located deep in the brain and are therefore difficult to reach. In the forebrain, they are also connected to parts of the hypothalamus, which controls the switch in basic behaviour, such as fight or flight and rest or sleep.

The findings of the team of researchers from the University of Freiburg reveal that certain intense sensory stimuli may affect such basic

behaviour through the activity of dopaminergic nerve cells. Because there are also connections between these nerve cells and the sensory organs, it is possible that dopaminergic nerve cells are involved in adjusting the sensitivity of sensory organs' reactions to stimuli. This function could be useful for treating diseases. The properties of dopaminergic [nerve cells](#) in the forebrain could thus be used in the future to reduce the sensation of patients with [restless legs syndrome](#) and hence to suppress the tingling in their extremities that occurs when sleeping. Further research of these dopaminergic neurons is expected to help scientists understand how these diseases develop – and in general how humans adapt to quickly changing stimuli and sensations in their environments.

More information: Sebastian Reinig et al. The Descending Diencephalic Dopamine System Is Tuned to Sensory Stimuli, *Current Biology* (2017). [DOI: 10.1016/j.cub.2016.11.059](https://doi.org/10.1016/j.cub.2016.11.059)

Provided by Albert Ludwigs University of Freiburg

Citation: Sensory stimuli control dopamine in the brain (2017, January 13) retrieved 26 December 2023 from <https://medicalxpress.com/news/2017-01-sensory-stimuli-dopamine-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--