

Both too much, too little weight tied to migraine

April 12 2017



Credit: Sasha Wolff/Wikipedia

Both obesity and being underweight are associated with an increased risk for migraine, according to a meta-analysis published in the April 12,

2017, online issue of *Neurology*, the medical journal of the American Academy of Neurology. The researchers looked at all available studies on body mass index (BMI) and migraine.

"As obesity and being underweight are potentially modifiable [risk factors](#) for migraine, awareness of these risk factors is vital for both people with migraine and doctors," said study author B. Lee Peterlin, DO, of Johns Hopkins University School of Medicine and a member of the American Academy of Neurology. "More research is needed to determine whether efforts to help people lose or gain weight could lower their risk for migraine."

A total of 12 studies with 288,981 participants were included in the meta-analysis. When the researchers compiled all of the results and adjusted for age and sex, they found that obese people were 27 percent more likely to have migraine than people of [normal weight](#). People who were underweight were 13 percent more likely to have migraine than people of normal weight.

Obesity was defined as a BMI of 30 or higher. Underweight was defined as a BMI of less than 18.5.

Peterlin said the risk between obesity and migraine was moderate and similar in size to the link between migraine and bipolar disorders and [ischemic heart disease](#), a condition of recurring chest pain or discomfort when part of the heart does not receive enough blood.

According to Peterlin, age and sex were important variables in the relationship between body mass index and migraine. "This makes sense, as the risk entailed by obesity and the risk of migraine is different in women and men and in younger and older people," she said. "Both obesity disease risk and the occurrence of migraine is more common in women and in younger people."

She continued, "It's not clear how [body composition](#) could affect migraine. Adipose tissue, or fatty tissue, secretes a wide range of molecules that could play a role in developing or triggering migraine. It's also possible that other factors such as changes in physical activity, medications, or other conditions such as depression play a role in the relationship between migraine and body composition."

Limitations of the meta-analysis include that for half of the studies people self-reported that they had [migraine](#) and for more than half of the studies [people](#) self-reported their [body mass index](#).

Provided by American Academy of Neurology

Citation: Both too much, too little weight tied to migraine (2017, April 12) retrieved 4 July 2023 from <https://medicalxpress.com/news/2017-04-weight-tied-migraine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.