

# Looking at the way we walk can help predict cognitive decline

October 28 2019

---



Credit: CC0 Public Domain

The way people walk is an indicator of how much their brains, as well as

their bodies, are aging. Scientists reporting in a special supplement to the *Journal of Alzheimer's Disease* (JAD) say that gait disorders, particularly slowing gait, should be considered a marker of future cognitive decline. They propose testing motor performance as well as cognitive performance in older adults with mild cognitive impairments.

"There is an emerging focus on the importance of assessing motor performance as well as [cognitive performance](#) to predict cognitive function loss," explained guest editor Manuel Montero-Odasso, MD, Ph.D., Departments of Medicine (Geriatric Medicine), and Epidemiology and Biostatistics, Schulich School of Medicine & Dentistry, University of Western Ontario, and Gait and Brain Lab, Parkwood Institute, Lawson Health Research Institute, London, ON, Canada. "In the past two decades, large epidemiological studies have shown that [gait disorders](#), particularly slowing [gait](#), may be present at early stages of dementia or may even predict who will be at risk of progressing to dementia. Subtle impairments in gait are more prevalent in [older adults](#) with cognitive impairments and dementia and are also associated with an increased risk of falls."

This supplement presents a comprehensive assessment of the current state of knowledge on the role of gait disturbances and quantitative gait analyses to be used as a motor biomarker to define subtypes of cognitive profiles and predict cognitive decline and dementia. It also evaluates screening tools that are important for developing strategies to prevent or delay onset of clinically meaningful change.

The issue showcases studies on:

- Epidemiology of gait disturbances and cognitive impairment
- Association between gait speed and further cognitive decline
- Structural brain volume covariance and its association with gait speed in patients with mild cognitive decline

- Association of cerebral amyloid-beta deposition and impaired [gait speed](#) and lower extremity function
- Use of the dual-task gait (DTG) paradigm of walking while performing a concurrent cognitively demanding task as a brain stress test to detect populations at risk
- Value of DTG to detect individuals at risk of cognitive decline in a cognitively healthy population of older adults
- Associations between DTG performance and cerebrospinal biomarkers of AD
- Effect of learning to use a mobility aid on gait and cognitive demands in AD—the use of assistive devices can help improve mobility and reduce the risk of falls but can also present a cognitive challenge
- Feasibility of measuring gait in an outpatient cognitive neurology setting



There's something in the way you walk. Credit: School of Medicine & Dentistry, University of Western Ontario, et al

Gait testing may help to detect the subgroup of at-risk patients who may benefit the most from invasive diagnostic procedures or early interventions. "We believe simple gait testing should be part of routine clinical assessment for older adults with cognitive impairments. Implementing this in clinics may be a challenge, but we hope the evidence presented in this issue will lead to progress in this area," noted guest editor George Perry, Ph.D., Editor-in-Chief of JAD, Professor of Biology, Semmes Distinguished University Chair in Neurobiology, The University of Texas at San Antonio, TX, USA.

"Finding early dementia detection methods is vital," added Dr. Montero-Odasso. "It is conceivable that in the future we will be able to make the diagnosis of AD and other dementias before people even have significant memory loss. In older adults with moderate [cognitive impairment](#), slowing down their usual walking by more than 20 percent when they add a cognitive task is indicative of a seven-fold increased risk to develop AD in a five-year timeframe. We believe that gait, as a complex brain-motor task, provides a golden window of opportunity to detect individuals at higher risk of dementia who can benefit the most from more invasive testing or early interventions."

Cognitive impairment and dementia carry a heavy healthcare burden. Around fifty million people worldwide have dementia. Every year there are nearly ten million new cases. AD is the most common form, accounting for around 60-70 percent of cases. Dementia is characterized by a progressive loss of cognitive function affecting memory, thinking, orientation, comprehension, calculation, learning capacity, language, and judgement. Impairments in gait are more common in [dementia](#) than in normal aging and may be related to the severity of cognitive decline.

**More information:** *Journal of Alzheimer's Disease* - Volume 71, issue s1: [content.iospress.com/journals/...eimers-disease/71/s1](http://content.iospress.com/journals/...eimers-disease/71/s1)

Provided by IOS Press

Citation: Looking at the way we walk can help predict cognitive decline (2019, October 28)  
retrieved 20 May 2023 from <https://medicalxpress.com/news/2019-10-cognitive-decline.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.