


Men more susceptible to COVID-19 than women

January 14 2021



Colorized scanning electron micrograph of a dying cell (blue) heavily infected with SARS-CoV-2 (yellow), the virus that causes COVID-19. Credit: NIAID Integrated Research Facility, Fort Detrick, Maryland.

Males are more likely to test positive for COVID-19, more likely to have complications and more likely to die from the virus than females, independent of age, according to a new study published this week in the

open-access journal *PLOS ONE* by Farhaan Vahidy of Houston Methodist Research Institute, US, and colleagues.

As the COVID-19 pandemic unfolds and evolves across the globe, researchers have identified population sub-groups with higher levels of disease vulnerability, such as those with advanced age or certain pre-existing conditions. Small studies from China and Europe have indicated that males tend to experience higher disease severity compared to [females](#). However a comprehensive analysis of COVID sex in a large and diverse US metropolitan area has been lacking.

In the new study, researchers used data from a large healthcare provider in the Houston, Texas [metropolitan area](#) to determine the associations between sex and COVID-19 epidemiology. Data on COVID testing, hospital stays, mortality and demographics were extracted from Electronic Medical Records (EMRs) of all 96,496 adults over 18 years old who were tested for SARS-CoV-2 by the health system between March 6 and August 22, 2020.

Overall, 15.5% (95% CI 15.3-15.8) of individuals in the cohort tested positive for SARS-CoV-2. After adjusting for sociodemographic factors and comorbidities, males had a higher likelihood of SARS-CoV-2 positivity (aOR 1.39, 95% CI 1.33-1.45) than females. Similarly, the proportion of patients requiring ICU care was significantly higher among males (34.1% (32.2—36.0)) as compared to females (27.6% (25.8—29.5)), OR: 1.36 (1.20—1.53). Moreover, more males (19.0% (17.5—20.6)) underwent [mechanical ventilation](#) than females (14.7% (13.3—16.2)), OR: 1.36 (1.17—1.59) and the proportion of [males](#) who experienced in-hospital mortality (11.6% (10.4—13.0)) was significantly higher as compared to females 8.3% (7.3—9.6), OR: 1.44 (1.18—1.75). The authors conclude that there is a clear and strong independent association between male sex and SARS-CoV-2 susceptibility, complications and poor outcomes and say that understanding sex

differences in the disease is a fundamental step toward improved disease management and intervention strategies for both men and women.

The authors add: "Males seem to be more likely to contract the SRAS-CoV-2 virus and also have a poor clinical course and outcomes related to COVID-19, compared to females. The exact contribution of gender and sex factors in susceptibility and outcomes of COVID-19 need further investigation."

More information: Farhaan S. Vahidy et al. Sex differences in susceptibility, severity, and outcomes of coronavirus disease 2019: Cross-sectional analysis from a diverse US metropolitan area, *PLOS ONE* (2021). [DOI: 10.1371/journal.pone.0245556](https://doi.org/10.1371/journal.pone.0245556)

Provided by Houston Methodist

Citation: Men more susceptible to COVID-19 than women (2021, January 14) retrieved 20 May 2023 from <https://medicalxpress.com/news/2021-01-men-susceptible-covid-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.