



Susceptibility of COPD patients to heart rate difference associated with exposure to metals in PM2.5

June 8 2021



Metal in PM2.5 exposure and difference in Heart rate in COPD and non-COPD participants. Credit: ©Science China Press

Epidemiological and toxicological studies indicate that the adverse outcomes of PM2.5 exposure associated closely with the chemical composition in PM2.5. Metals in PM2.5 are highly concerned for their induced disruption of iron homeostasis in the lung and following oxidative stress, which is one of the key mechanisms underlying the

cardiovascular autonomic dysfunction of PM_{2.5} exposure. However, there is no clear evidence on whether COPD patients are more susceptible to cardiovascular autonomic dysfunction associated with exposure to metals in ambient PM_{2.5} than individuals without COPD. Based on a panel study, the researchers directly compared metal-associated cardiovascular autonomic dysfunction between COPD patients and healthy controls.

"We observed higher levels of heart rate (HR) associated with exposure to PM_{2.5} and the [metal](#) elements therein. Associations between HR and metals exposure in COPD patients were stronger than those in healthy controls. Exposure to Cr had robust associations with HR, comparing to other metals." said Dr. Ke Gao, the first author for this work.

This study has many advantages. "It is a panel study design with repeated measurements of HR and exposure to metals in PM_{2.5}. All of the recruited participants acted as their own controls, thereby lessening deviation linked to intra-[individual differences](#). In addition, we recruited both individuals with COPD and healthy controls, allowing them to make direct comparisons of different responses from the COPD and healthy groups and to investigate the susceptibility of individuals with COPD to air pollution-linked responses." said Dr. Xi Chen, the co-first author for this work.

"Although our results require further validation, they shed light on the disease burden of COPD associated with exposure to metals in airborne PM_{2.5} in China." Said Prof. Tong Zhu, the corresponding author of the paper.

These encouraging results provide valuable data and improve the scientific understanding of susceptibility of COPD patients to metal-associated cardiovascular autonomic dysfunction.

More information: Ke Gao et al, Susceptibility of patients with chronic obstructive pulmonary disease to heart rate difference associated with the short-term exposure to metals in ambient fine particles: A panel study in Beijing, China, *Science China Life Sciences* (2021). [DOI: 10.1007/s11427-020-1912-4](https://doi.org/10.1007/s11427-020-1912-4)

Provided by Science China Press

Citation: Susceptibility of COPD patients to heart rate difference associated with exposure to metals in PM_{2.5} (2021, June 8) retrieved 21 July 2023 from <https://medicalxpress.com/news/2021-06-susceptibility-copd-patients-heart-difference.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.