

# Vitamin D may protect against young-onset colorectal cancer

August 17 2021

---



Cancer — Histopathologic image of colonic carcinoid. Credit: Wikipedia/CC BY-SA 3.0

Consuming higher amounts of Vitamin D—mainly from dietary sources—may help protect against developing young-onset colorectal cancer or precancerous colon polyps, according to the first study to show such an association.

The study, recently published online in the journal *Gastroenterology*, by scientists from Dana-Farber Cancer Institute, the Harvard T.H. Chan School of Public Health, and other institutions, could potentially lead to recommendations for higher [vitamin D](#) intake as an inexpensive complement to screening tests as a colorectal [cancer](#) prevention strategy for adults younger than age 50.

While the overall incidence of colorectal cancer has been declining, cases have been increasing in younger adults—a worrisome trend that has yet to be explained. The authors of the study, including senior co-authors Kimmie Ng, MD, MPH, of Dana-Farber, and Edward Giovannucci, MD, DSc., of the T.H. Chan School, noted that vitamin D intake from food sources such as fish, mushrooms, eggs, and milk has decreased in the past several decades. There is growing evidence of an association between vitamin D and risk of colorectal cancer mortality. However, prior to the current study, no research has examined whether total vitamin D intake is associated with the risk of young-onset colorectal cancer.

"Vitamin D has known activity against colorectal cancer in laboratory studies. Because vitamin D deficiency has been steadily increasing over the past few years, we wondered whether this could be contributing to the rising rates of colorectal cancer in young individuals," said Ng, director of the Young-Onset Colorectal Cancer Center at Dana-Farber. "We found that total vitamin D intake of 300 IU per day or more—roughly equivalent to three 8-oz. glasses of milk—was associated with an approximately 50% lower risk of developing young-onset colorectal cancer."

The results of the study were obtained by calculating the total vitamin D intake—both from dietary sources and supplements—of 94,205 women participating in the Nurses' Health Study II (NHS II). This study is a prospective cohort study of nurses aged 25 to 42 years that began in

1989. The women are followed every two years by questionnaires on demographics, diet and lifestyle factors, and medical and other health-related information. The researchers focused on a primary endpoint—young-onset colorectal cancer, diagnosed before 50 years of age. They also asked on a follow-up questionnaire whether they had had a colonoscopy or sigmoidoscopy where colorectal polyps (which may be precursors to colorectal cancer) were found.

During the period from 1991 to 2015 the researchers documented 111 cases of young-onset colorectal cancer and 3,317 colorectal polyps. Analysis showed that higher total vitamin D intake was associated with a significantly reduced risk of early-onset colorectal cancer. The same link was found between higher vitamin D intake and risk of colon polyps detected before age 50.

The association was stronger for dietary vitamin D—principally from dairy products—than from vitamin D supplements. The study authors said that finding could be due to chance or to unknown factors that are not yet understood.

Interestingly, the researchers didn't find a significant association between total vitamin D intake and risk of colorectal cancer diagnosed after age 50. The findings were not able to explain this inconsistency, and the scientists said further research in a larger sample is necessary to determine if the protective effect of vitamin D is actually stronger in young-onset colorectal cancer.

In any case, the investigators concluded that higher total vitamin D intake is associated with decreased risks of young-onset colorectal cancer and precursors (polyps). "Our results further support that vitamin D may be important in younger adults for health and possibly colorectal cancer prevention," said Ng. "It is critical to understand the risk factors that are associated with young-onset colorectal cancer so that we can

make informed recommendations about diet and lifestyle, as well as identify high risk individuals to target for earlier screening."

The study was funded by grants from the U.S. National Institutes of Health and the Department of Defense; by the American Cancer Society Mentored Research Scholar Grant; and by the Project P Fund.

Ng's disclosures include research funding from Pharmavite, Revolution Medicines, Janssen, and Evergrande Group; Advisory boards for Array Biopharma, Seattle Genetics, and BiomX; and consulting for X-Biotix Therapeutics.

**More information:** Hanseul Kim et al, Total Vitamin D Intake and Risks of Early-Onset Colorectal Cancer and Precursors, *Gastroenterology* (2021). [DOI: 10.1053/j.gastro.2021.07.002](https://doi.org/10.1053/j.gastro.2021.07.002)

Provided by Dana-Farber Cancer Institute

Citation: Vitamin D may protect against young-onset colorectal cancer (2021, August 17) retrieved 9 April 2023 from <https://medicalxpress.com/news/2021-08-vitamin-d-young-onset-colorectal-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--