

Childhood asthma rates stable in Western countries but increasing in lower-middle income countries

October 29 2021



Credit: CC0 Public Domain

The prevalence of asthma in children was found to have decreased in low-income countries, but increased in lower-middle income countries,

according to the latest phase of the largest face-to-face health survey ever conducted, published in *The Lancet*.

The prevalence in upper-middle and high-income countries was stable, found the Global Asthma Network (GAN) study, which shows how the worldwide burden of children's [asthma](#) symptoms is changing.

Led by the University of Auckland with partners including the London School of Hygiene & Tropical Medicine (LSHTM), the Phase 1 study included nearly 120,000 children and adolescents from 27 centers in 14 countries, and continues a series of global surveys over 27 years (between 1993 to 2020).

Three phases of data collection have overall included more than 300 centers in more than 100 countries. The first two phases were conducted by the International Study of Asthma and Allergies in Childhood (ISAAC), which has now continued as the Global Asthma Network.

Asthma is the most common non-communicable disease in children, representing a large burden of disease globally with about 10-15% of children globally having had [asthma symptoms](#) in the last year. The overall prevalence in the centers that took part in the new survey had changed little in the last 15-20 years. Prevalence increased by about 1% every 10 years in Europe, and by more than 2% in Africa and the Middle East, whereas prevalence had fallen by about 1% in Asia-Pacific, and was unchanged in the Americas.

LSHTM led the latest round of data collection using a survey of self-reported asthma symptoms, and has hosted one of the two data centers. The report includes new data showing time trends in asthma symptoms for two age groups of children: 6-7 year olds and 13-14 year olds.

This study provides important clues to what may be causing the global

changes in asthma prevalence, and what we can do to prevent it.

Professor Neil Pearce from LSHTM, Coordinator of the latest round of data collection, said: "These results suggest that asthma is a disease of westernization, with rates increasing in some low- and middle-income countries as they become more westernized. There is an urgent need for improved asthma management globally, and better access to inexpensive asthma medicines.

"Currently many children with asthma don't get diagnosed, not only on low-and-middle-income countries, but also in the United Kingdom. This study, among other surveys, is the only way to get accurate data."

The researchers acknowledge the limitations of this study, include the small number of centers involved in this phase, which makes it difficult to generalize the findings to global prevalence changes.

More information: M Innes Asher et al, Worldwide trends in the burden of asthma symptoms in school-aged children: Global Asthma Network Phase I cross-sectional study, *The Lancet* (2021). [DOI: 10.1016/S0140-6736\(21\)01450-1](https://doi.org/10.1016/S0140-6736(21)01450-1)

Provided by London School of Hygiene & Tropical Medicine

Citation: Childhood asthma rates stable in Western countries but increasing in lower-middle income countries (2021, October 29) retrieved 20 November 2023 from <https://medicalxpress.com/news/2021-10-childhood-asthma-stable-western-countries.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.