


PTSD symptoms vary over course of menstrual cycle

October 28 2021



Credit: Unsplash/CC0 Public Domain

In women who have experienced trauma, post-traumatic stress disorder symptoms may vary over the course of the menstrual cycle, with more symptoms during the first few days of the cycle when the hormone

estradiol is low, and fewer symptoms close to ovulation, when estradiol is high, finds research published by the American Psychological Association.

The results could have implications for PTSD diagnosis and treatment, according to lead author Jenna Rieder, Ph.D., an assistant professor of psychology at Thomas Jefferson University in Philadelphia. "When in the [cycle](#) you assess women might actually affect whether they meet diagnostic criteria for PTSD, especially for people who are right on the border," she said. "And that can have real practical implications, say for someone who is a veteran and entitled to benefits or for health insurance purposes."

The research was published in the journal *Psychological Trauma: Theory, Research, Practice and Policy*.

Estradiol is a form of estrogen that regulates the [reproductive cycle](#) in women. During the follicular phase of the menstrual cycle, rising [estradiol levels](#) trigger a cascade of events that result in ovulation. Studies have linked low-estradiol portions of the cycle to greater activation in the limbic areas of the brain, which are related to emotion, and to lower activation in the prefrontal cortex when viewing emotional content. Low estradiol has also been linked to greater stress and anxiety as well as increased [fear responses](#).

In order to examine whether those links might affect trauma response, researchers studied 40 women, ages 18 to 33, all of whom had experienced or witnessed a traumatic event, such as a serious injury or sexual violence. In the first part of the study, which took place in a research lab, researchers measured the participants' level of estradiol in their saliva, then asked them to describe the trauma that had happened to them and the PTSD symptoms they'd experienced in the past month. They found that lower estradiol was associated with greater self-reported

[symptom](#) severity in the participants.

The researchers also measured two stress biomarkers in participants' saliva, the hormone cortisol and the enzyme salivary alpha-amylase, before and after the participants described their trauma. Salivary alpha-amylase is related to the "fight-or-flight" stress response and cortisol is related to the body's slower, more sustained stress response.

"In a healthy system we want a moderate, coordinated response of both of these biomarkers," Rieder said. In the women in the low-estradiol portions of their menstrual cycles, the researchers instead found low cortisol and high salivary alpha-amylase levels in response to retelling their trauma stories—a pattern that's been linked in previous studies with maladaptive stress responses.

In the second part of the study, the researchers asked the participants to answer five daily questionnaires (upon waking, before bed, and at three times during the day), for 10 days spanning the high- and low-estradiol portions of their menstrual cycles. The questionnaires measured how participants were feeling at each time (from "extremely unpleasant" to "extremely pleasant," and "extremely non-stimulated or activated" to "extremely stimulated or activated"). Participants also completed a PTSD symptom checklist each evening.

On average, the researchers found that participants had greater variability in their daily moods during the low-[estradiol](#) days of their cycle and reported more severe PTSD symptoms on those days.

The findings could have implications for diagnosing and treating PTSD in women, who have long been underrepresented in PTSD research. "PTSD for a long time was mostly studied in men, in part because it was mainly studied in veterans, who were mostly men," Rieder said.

In addition to affecting diagnosis, knowing how the menstrual cycle affects PTSD symptoms could be useful for both clinicians and patients, according to Rieder. "I think this is something that clinicians would want to know, so they can impart this knowledge as part of psychoeducation," Rieder said. "For [women](#) who are naturally cycling, it may be useful to understand how the menstrual cycle affects their symptoms. When you can explain what's happening biologically, it often becomes less threatening."

More information: "Estradiol, Stress Reactivity, and Daily Affective Experiences in Trauma-Exposed Women," *Psychological Trauma: Theory, Research, Practice and Policy* [DOI: 10.1037/tra0001113](https://doi.org/10.1037/tra0001113)

Provided by American Psychological Association

Citation: PTSD symptoms vary over course of menstrual cycle (2021, October 28) retrieved 31 January 2024 from <https://medicalxpress.com/news/2021-10-ptsd-symptoms-vary-menstrual.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--