


Reducing tau protein levels can help maintain balance in the brain, says study

October 19 2021, by Julie Langelier



Credit: CC0 Public Domain

With the school year well on its way, we all recognize the importance of getting students energized. However, we also know that overstimulated



children can wreak havoc on a classroom, leaving a teacher trying to regain control of their group. The situation is very similar in the brain, where cells must remain active—but in an orderly manner—to carry out their many important tasks.

If too many excitatory <u>cells</u> (the students in this metaphor) begin firing at the same time, the overexcitation can spread through an entire neural network (the classroom), resulting in epileptic activity. So, inhibitory cells (the teachers) tamp down the excitation to maintain balance in the brain.

Now, researchers at Gladstone Institutes have uncovered that reducing levels of the protein tau, which is known for its role in Alzheimer's disease and other neurodegenerative conditions, changes excitatory and inhibitory cells in ways that make it harder for the brain to burst with overexcitation.

"Epilepsy can be associated with many brain disorders, including Alzheimer's disease and autism," says Lennart Mucke, MD, director of the Gladstone Institute of Neurological Disease and senior author of the new study published in the journal *Cell Reports*. "We previously showed in mouse models that reducing tau levels makes the brain more resistant to epilepsy of diverse causes, but we have puzzled about the underlying mechanisms. Our recent findings shed light onto how tau reduction affects different types of brain cells involved in creating abnormal network activity."

It is critical to preserve a balance between the activity of excitatory and inhibitory cells for the brain to process information, appropriately control movements, and perceive things as they really are, rather than in a distorted way.

"The diseases we're interested in have in common that they cause an



excitation-inhibition imbalance in the brain," says Mucke, who is also the Joseph B. Martin Distinguished Professor of Neuroscience and a professor of neurology at UC San Francisco.

Mucke and his team discovered that reducing levels of tau impacts both excitatory and <u>inhibitory cells</u>, but in different ways. They showed that excitatory neurons that lack tau don't fire as much, whereas inhibitory neurons that lack tau are more easily stimulated to fire.

Together, these effects reduce the excitation-inhibition ratio in brain networks, counteracting diseases that cause abnormal increases in that ratio. To return to the classroom example, the neural effects of reducing tau are the equivalent of making the students less rambunctious and the teachers more effective at supervising them.

Just like calm in the classroom facilitates learning, the calming effect of tau reduction on brain networks might improve the brain's ability to perform its functions, especially when the calm is being disrupted by <u>disease</u>. The insights gained in the new study could help guide the development of novel therapies that target tau as a way of reducing abnormal <u>brain</u> activities in a range of neurological and psychiatric diseases.

More information: Lennart Mucke, Tau reduction affects excitatory and inhibitory neurons differentially, reducing excitation/inhibition ratios and counteracting network hypersynchrony, *Cell Reports* (2021). DOI: 10.1016/j.celrep.2021.109855. www.cell.com/cell-reports/full ... 2211-1247(21)01322-X

Provided by Gladstone Institutes



Citation: Reducing tau protein levels can help maintain balance in the brain, says study (2021, October 19) retrieved 20 November 2023 from <u>https://medicalxpress.com/news/2021-10-tau-protein-brain.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.