

Increased take-home methadone during pandemic did not worsen outcomes

November 1 2021



Buprenorphine. Credit: Pixabay/CC0 Public Domain

Relaxing limits on take-home doses of methadone—a medication used to treat opioid addiction—does not appear to lead to worse treatment outcomes, according to a new study led by Washington State University

researchers.

Published in the *American Journal of Drug and Alcohol Abuse*, the study looked at the impact of a temporary policy change allowing providers to send patients home with additional [methadone](#) doses during the COVID-19 pandemic. Previously, federal regulations allowed take-home privileges only for established patients who have proven themselves stable, a measure intended to reduce risk of patients selling the provided methadone.

While the researchers saw the average number of methadone take-home doses nearly double during the pandemic, they found no significant changes in treatment outcomes. There was no rise in the number of patients experiencing emergency department visits, whether overdose-related or for other reasons. There was also no notable increase in the number of patients who tested negative for methadone.

"Our research highlights the need to consider permanently loosening the restrictions on methadone take-home doses, which would help many people who are struggling to access opioid treatment," said lead study author Ofer Amram, an assistant professor in the WSU Elson S. Floyd College of Medicine whose research focuses on decreasing health disparities in vulnerable populations.

The study was based on data from 183 patients treated at a methadone clinic in Spokane County, Washington. Methadone take-home doses from the clinic rose from 11.4 doses per 30 days in the eight months prior to the pandemic policy change to 22.3 doses per 30 days in the eight months following.

Methadone can only be prescribed and dispensed by federally approved opioid treatment programs, and there are only about 1,800 such programs around the country. This makes requiring daily visits to get

methadone especially difficult for individuals who live far from a treatment program, Amram said.

In a previously published study of patients attending the same clinic involved in this study—the only publicly funded methadone clinic in Eastern Washington—Amram and his coauthors showed that patients who lived closer to the clinic continued with treatment better than those who lived further away. Yet take-home limitations pose challenges even for those who live close to a [treatment](#) clinic but don't have reliable access to transportation or find that daily visits take too much time away from employment, Amram said.

More information: Ofer Amram et al, The impact of relaxation of methadone take-home protocols on treatment outcomes in the COVID-19 era, *The American Journal of Drug and Alcohol Abuse* (2021). [DOI: 10.1080/00952990.2021.1979991](https://doi.org/10.1080/00952990.2021.1979991)

Provided by Washington State University

Citation: Increased take-home methadone during pandemic did not worsen outcomes (2021, November 1) retrieved 24 March 2023 from <https://medicalxpress.com/news/2021-11-take-home-methadone-pandemic-worsen-outcomes.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|