

Big differences in how emergency doctors assess patient care needs

December 2 2021, by Svein Inge Meland



Credit: Pixabay/CC0 Public Domain

When researchers divided out-of-hours general practitioners (GPs) into four groups based on how often they admit their patients to the hospital, they found that if you are seen by a doctor who belongs to the 25 per

cent who admit the most often, your chance of a hospital stay is much greater.

Patients over the age of 69 have almost double the chance of being admitted to hospital by GPs with [higher admission rates](#) as compared with patients who see a GP in the lowest [admission](#) rate group.

One might reasonably believe that the patients who have seen the most restrictive out-of-hours (emergency) doctors will eventually have more hospital admissions down the road. To some extent this does happen.

However, 30 days after an emergency room visit, findings still show fewer admissions among the patients who were seen by the most restrictive GPs than among those whose physicians were most likely to send the patient to hospital.

Expensive for hospitals

"This means that the GPs with the lowest threshold for admission inflict significant costs on the hospitals that might be unnecessary," says Ellen Rabben Svedahl, a Ph.D. candidate at the Norwegian University of Science and Technology (NTNU) and first author of the study.

"Our study findings show that a GPs' out-of-hours decision to admit a patient affects the total use of [health services](#) and hospital costs. We've calculated that the doctors who admit patients the most often cost about NOK 4.2 million more per 1000 emergency room visits than the GP group with the fewest referrals," Svedahl said.

With approximately 200 000 emergency consultations a year for the over-69 age group, this becomes a major cost. Svedahl and her supervisor Johan Håkon Bjørngaard, a professor in the Department of Public Health and Nursing, are part of a group at NTNU that studies

health services, funded by the Research Council of Norway.

Risk of death not any higher

But doesn't the risk to patients' lives and health increase when they're seen by doctors who have the highest threshold for admission?

Svedahl says the researchers didn't find that different admission thresholds led to differences in patient mortality over the 30 days following the out-of-hours visit.

"At the same time, we know that too few—or too many—hospitalizations can potentially mean an increased risk of complications, especially for the elderly. What we didn't investigate is how satisfied the patients were, or whether any complications arose in the long term," Svedahl said.

Gender and age differences

The researchers grouped doctors according to their frequency of hospital admissions, independent of gender and age. But they also found that men and women, as well as younger and older doctors, have different practices. When children aged 0-10 years are seen by a female GP in the emergency room, the admission rate is 25 per cent higher than if the GP were a man.

The propensity to admit patients decreases by 5-8 per cent with every decade older that a physician is. Specialist-trained doctors are a little less likely to refer their patients to hospitals, while GPs with low out-of-hours experience in the previous two years tend to admit more.

Riskier to see a male doctor?

Gender and age differences have been found in previous studies, but little research has been done on the consequences of different thresholds for admission.

"Although the doctor's gender and age indicate some differences, they're quite small and even out somewhat over time after the out-of-hours contact. The costs for the specialist health services thus aren't that much lower, even though male and older doctors are more restrictive in referring patients to hospital," says Svedahl.

Slightly higher mortality

The researchers found that mortality was slightly higher (12-17 per cent) among patients seen by doctors who had less emergency room practice during the previous two years.

"Here we have to allow for some uncertainty. However, we can't ignore the fact that some patients are sent home who should have been hospitalized. Another possible explanation is that older, more fragile people were admitted who shouldn't have been sent to hospital. This should be further investigated," Svedahl says.

Warns against raising the threshold

Does that mean that doctors should raise the threshold for hospitalization?"

"The question is rather how we can ensure the quality of admissions in the best possible way. The study shows that a more restrictive hospitalization practice could affect patient safety," says Svedahl.

In Norway, GPs are the largest group of physicians that rotate through emergency room shifts. Limited time and little knowledge of the patients

means that out-of-hours doctors have a poorer basis for making their assessments than in their primary care practice as a GP.

Svedahl believes better systems—such as greater opportunities to confer with another doctor or specialist, introduction of shared patient records and more systematic feedback on how patients are doing after referral—could contribute to increasing the quality of care.

Admission in emergency care centers

Some municipalities have established out-of-hours emergency care centers where patients can be admitted.

"Schemes like this give the emergency doctor several options. Instead of admitting patients with questionable needs to hospitals or sending them home, they can be referred to a 24-hour unit in the out-of-hours primary care centers where they receive treatment, care and are observed by GPs and nurses," says Svedahl.

An aging population and higher life expectancy mean that the need for emergency care and specialist treatment will increase. Svedahl, who is also a physician, is concerned that the work pressure among GPs could weaken out-of-hours services.

"Many GPs report high work pressure. This could mean that we'll have fewer doctors with a lot of experience in out-of-hours emergency centers. This situation might in turn lead to more [hospital](#) admissions, higher costs and increased pressure on the specialist health services," she says.

Careful study design

Studying hospitalization practices is difficult due to varying morbidity in

the doctors' patient groups.

"We were careful about how we made the comparisons. A patient in his 70s who saw a doctor in the emergency room on a Wednesday was only compared with other patients in their 70s who saw another doctor in the same emergency room on the other Wednesdays in the same month. This means that we only compared patients who saw other doctors in the same emergency room for comparable periods of time. With this design, we assumed that the differences we found would not be due to the different doctors having [patients](#) with varying morbidity," Svedahl explains.

The researchers used data from all [emergency room visits](#) in the period 2008-2016.

Seeing how services are linked

"The study is another example that points out the need for primary and specialist health services to be seen in a linked context," says Johan Håkon Bjørngaard, the senior author on the article.

"An intensified effort in out-of-hours emergency care is more expensive for the municipalities, but can be better for the patient. It can also reduce the pressure on the specialist health services and would be a good use of resources for society," he said.

The research was published in *Family Practice*.

More information: Ellen Rabben Svedahl et al, Effects of GP characteristics on unplanned hospital admissions and patient safety. A 9-year follow-up of all Norwegian out-of-hours contacts, *Family Practice* (2021). [DOI: 10.1093/fampra/cmab120](https://doi.org/10.1093/fampra/cmab120)

Provided by Norwegian University of Science and Technology

Citation: Big differences in how emergency doctors assess patient care needs (2021, December 2) retrieved 20 July 2023 from

<https://medicalxpress.com/news/2021-12-big-differences-emergency-doctors-patient.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.