

# Undiagnosed autistic traits common among youths with substance use disorders

January 24 2022

---



Credit: Unsplash/CC0 Public Domain

One in five teens and young adults who seek treatment for alcohol or drug use may have traits characteristic of a previously unrecognized autism spectrum disorder (ASD), researchers at Massachusetts General

Hospital (MGH) report.

They found that among patients with an average age of 18.7 years being treated in an outpatient [substance use disorder](#) (SUD) clinic, 20% had elevated scores on the Social Responsiveness Scale-2 (SRS-2), a parent- or teacher-reported measure that has been shown to reliably identify the presence and severity of social impairment among individuals along the autism spectrum, and to distinguish autism from other [disorders](#).

The study, results of which are published in *The American Journal on Addictions*, is the first to look at the prevalence of previously undiagnosed autistic traits among teens and [young adults](#) with SUD, says lead author James McKowen, Ph.D., of the Addiction Recovery Management Service at MGH and Harvard Medical School.

"Usually studies of substance use disorder in autism are done in those with an autism diagnosis already," he says. "We have looked at this question from the other side, asking how many people with substance use disorder have autism."

The researchers asked parents of 69 youths reporting for the first time to a specialty outpatient psychiatric SUD clinic to fill out the SRS-2 form. The form is designed to measure an individual's social awareness, social cognition (thinking about other people and interactions with them), social communication, social motivation, and restricted interests and repetitive behaviors.

They found that although there were few differences between those with elevated autistic trait scores and those with lower, non-autistic scores in terms of demographic or psychiatric factors, the adolescents with higher SRS-2 scores had a nearly eightfold higher likelihood of stimulant use disorder, and a fivefold higher risk for opioid use disorder.

The findings highlight the importance of assessing patients in a SUD treatment setting for autistic traits, the researchers write.

"For clinicians, the big takeaway point from this study is that we need to get better at screening and certainly training in the presence of autism spectrum disorder, because many clinicians treat substance use disorder but don't have specialty developmental training, particularly for issues around [autism](#)," McKowen says.

"For parents, the big takeaway is that if you suspect that your child may have an [autism spectrum](#) issue or if school personnel have suggested that your child may have autistic traits, you should certainly get that assessed, and let your clinicians know whether your child has had a prior diagnosis of ASD," he says.

The researchers are developing a free clinical therapy protocol that can help clinicians better address the issues of autistic traits in patients with SUD.

**More information:** James McKowen et al, Characterizing autistic traits in treatment-seeking young adults with substance use disorders, *The American Journal on Addictions* (2021). [DOI: 10.1111/ajad.13247](https://doi.org/10.1111/ajad.13247)

Provided by Massachusetts General Hospital

Citation: Undiagnosed autistic traits common among youths with substance use disorders (2022, January 24) retrieved 9 April 2023 from <https://medicalxpress.com/news/2022-01-undiagnosed-autistic-traits-common-youths.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.