

One in five male adolescents suffers from high blood pressure

May 13 2022, by Johannes Angerer



Credit: Pixabay/CC0 Public Domain

Unhealthy levels of hypertension are increasing globally, especially during puberty, with boys affected three to four times more frequently than girls: Around 20% of male adolescents have elevated blood pressure. The main causes of primary hypertension (hypertension not triggered by another disease) in young people are obesity and lack of exercise, and increasingly, chronic psychological stress.

Based on [scientific data](#), a main reason for the increasing health risk in [young people](#) is obesity: while primary hypertension is found in 1.4% of normal-weight adolescents and 7.1% of overweight adolescents, the percentage rises to 25% in obese adolescents. "Increased waist circumference presenting the abdominal fat in particular, is associated with high [blood pressure](#) and early onset of cardiovascular diseases," reports Susanne Greber-Platzer, Head of the Department of Pediatrics and Adolescent Medicine at MedUni Vienna.

Anxiety, pressure and stress

Lack of exercise, in turn, leads to a threefold increase in the risk of hypertension. Chronic exposure to anxiety, pressure and stress is also increasing in children and adolescents, with significant effects on their health and blood pressure. Additionally [sugary drinks](#) and salty foods are directly affecting the blood pressure. Rapid growing during puberty, especially in boys, often leads to pathological blood pressure values. Prematurity, low birth weight and genetic predisposition may also play a role.

Factors such as age, gender and height must be taken into account when measuring hypertension in children under 15 years of age. For

adolescents aged 16 and older, a blood pressure above 140/90 is considered as hypertensive after three independent measurements, the same as in adults. Like adults, children and adolescents do not notice [high blood pressure](#), therefore early diagnosis is challenging. General symptoms like headache, dizziness or nose bleeding can occur, but are not necessarily present. Hypertensive crises, markedly elevated systolic blood pressure values, can lead to confusion and seizures and even unconsciousness, presenting life threatening situations.

High risk for early onset of cardiovascular diseases

Long-term high blood pressure leads to left [ventricular hypertrophy](#) and endothelial damage of the arterial vessels including coronary arteries, consequently increasing the risk for premature myocardial infarction or stroke, even in young adulthood. "To prevent the risk of early cardiovascular damage, a screening from the age of three years would be necessary," says Susanne Greber-Platzer, emphasizing the importance of measuring blood pressure in children and adolescents at least every two years. Therefore, a [healthy lifestyle](#) should be forced and additional medication to lower the blood pressure can be indicated in childhood to prevent early cardiovascular damage.

If hypertension is caused by other diseases (secondary hypertension), it is necessary to treat the primary disease appropriately and enabling blood pressure reduction. For example, Diabetes mellitus type I causes high blood pressure in about 15%, which frequently manifests in childhood or adolescence.

Provided by Medical University of Vienna

Citation: One in five male adolescents suffers from high blood pressure (2022, May 13) retrieved 14 July 2023 from <https://medicalxpress.com/news/2022-05-male-adolescents-high-blood->

[pressure.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.