


Recurrent UTIs linked to gut microbiome, chronic inflammation

May 3 2022, by Tamara Bhandari





Study design and sample collection for the UMB study. Credit: *Nature Microbiology* (2022). DOI: 10.1038/s41564-022-01107-x



One of the greatest frustrations regarding urinary tract infections (UTIs) is that they so often recur. UTIs are caused by bacteria in the urinary tract and characterized by frequent and painful urination. A round of antibiotics usually clears up the symptoms, but the relief is often temporary: A quarter of women go on to develop a second UTI within six months. Some unfortunate individuals get UTIs over and over, and require antibiotics every few months.

A new study suggests that women who get recurrent UTIs may be caught in a <u>vicious cycle</u> in which antibiotics given to eradicate one infection predispose them to develop another. The study, by researchers at Washington University School of Medicine in St. Louis and the Broad Institute of MIT and Harvard, showed that a round of antibiotics eliminates disease-causing <u>bacteria</u> from the bladder but not from the intestines. Surviving bacteria in the gut can multiply and spread to the bladder again, causing another UTI.

At the same time, repeated cycles of antibiotics wreak havoc on the community of helpful bacteria that normally live in the intestines, the so-called gut microbiome. Similar to other disorders in which gut microbes and the immune system are linked, women with recurrent UTIs in the study had less diverse microbiomes that were deficient in an important group of bacteria that helps regulate inflammation, and a distinct immunological signature in their blood indicative of inflammation.

The study is published May 2 in *Nature Microbiology*.

"It's frustrating for women who are coming in to the doctor with recurrence after recurrence after recurrence, and the doctor, who's typically male, gives them advice about hygiene," said co-senior author Scott J. Hultgren, the Helen L. Stoever Professor of Molecular Microbiology at Washington University. "That's not necessarily what the problem is. It's not necessarily poor hygiene that's causing this. The



problem lies in the disease itself, in this connection between the gut and the bladder and levels of inflammation. Basically, physicians don't know what to do with recurrent UTI. All they have is antibiotics, so they throw more antibiotics at the problem, which probably just makes things worse."

Most UTIs are caused by Escherichia coli (E. coli) bacteria from the intestines that get into the <u>urinary tract</u>. To understand why some women get infection after infection and others get one or none, Hultgren teamed up with Broad Institute scientists Ashlee Earl, the senior group leader for the Bacterial Genomics Group at Broad and the paper's co-senior author, and Colin Worby, a computational biologist and the paper's lead author.

The researchers studied 15 women with histories of recurrent UTIs and 16 women without. All participants provided urine and <u>blood samples</u> at the start of the study and monthly stool samples. The team analyzed the bacterial composition in the stool samples, tested the urine for the presence of bacteria, and measured gene expression in blood samples.

Over the course of a year, 24 UTIs occurred, all in participants with histories of repeated UTIs. When participants were diagnosed with a UTI, the team took additional urine, blood and stool samples.

The difference between the women who got repeated UTIs and those who didn't, surprisingly, didn't come down to the kind of E. coli in their intestines or even the presence of E. coli in their bladders. Both groups carried E. coli strains in their guts capable of causing UTIs, and such strains occasionally spread to their bladders.

The real difference was in the makeup of their gut microbiomes. Patients with repeat infections showed decreased diversity of healthy gut microbial species, which could provide more opportunities for disease-causing species to gain a foothold and multiply. Notably, the



microbiomes of women with recurrent UTIs were particularly scarce in bacteria that produce butyrate, a short-chain fatty acid with anti-inflammatory effects.

"We think that women in the control group were able to clear the bacteria from their bladders before they caused disease, and <u>women</u> with recurrent UTI were not, because of a distinct immune response to bacterial invasion of the bladder potentially mediated by the <u>gut microbiome</u>," Worby said.

The findings highlight the importance of finding alternatives to antibiotics for treating UTIs.

"Our study clearly demonstrates that antibiotics do not prevent future infections or clear UTI-causing strains from the gut, and they may even make recurrence more likely by keeping the microbiome in a disrupted state," Worby said

Hultgren has long worked on finding innovative therapies to eradicate disease-causing strains of E. coli from the body while sparing the rest of the bacterial community. His research forms the basis of an experimental drug based on the sugar mannoside and an investigational vaccine, both of which are being tested in people. Another strategy would be to rebalance the microbiome through fecal transplants, probiotic foods or other means.

"This is one of the most common infections in the United States, if not the world," Hultgren said. "A good percentage of these UTI patients go on to get these chronic recurrences, and it results in decreased quality of life. There is a real need to develop better therapeutics that break this vicious cycle."

More information: Scott Hultgren, Longitudinal multi-omics analyses



link gut microbiome dysbiosis with recurrent urinary tract infections in women, *Nature Microbiology* (2022). DOI: 10.1038/s41564-022-01107-x . www.nature.com/articles/s41564-022-01107-x

Provided by Washington University School of Medicine in St. Louis

Citation: Recurrent UTIs linked to gut microbiome, chronic inflammation (2022, May 3) retrieved 5 February 2023 from https://medicalxpress.com/news/2022-05-recurrent-utis-linked-gut-microbiome.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.