


Ultrasound-assisted laser technique vaporizes artery plaque

May 24 2022



Atherosclerosis, a buildup of plaque, can lead to heart disease, artery disease, and chronic kidney disease and is traditionally treated by inserting and inflating a balloon to expand the artery. Credit: Rohit Singh, University of Kansas

Atherosclerosis, a buildup of plaque, can lead to heart disease, artery disease, and chronic kidney disease and is traditionally treated by inserting and inflating a balloon to expand the artery. Other treatments based on lasers can remove blockages rather than simply compressing

them but are used infrequently, because they have a high risk of complication and low efficacy.

Rohit Singh, of the University of Kansas, and other researchers developed a method that combines a low-power [laser](#) with ultrasound to remove arterial plaque safely and efficiently. Singh will describe preliminary results in his presentation, "A novel ultrasound-assisted [laser technique](#) to remove [atherosclerotic plaques](#)," at the 182nd Meeting of the Acoustical Society of America. The session will take place May 24 at 5:05 p.m. Eastern U.S. at the Sheraton Denver Downtown Hotel.

High-power laser treatments direct [thermal energy](#) to vaporize water in the artery and create a vapor bubble, which expands and collapses to break the plaque. Similarly, the technology, pioneered by Xinmai Yang, doctoral advisor for the team, uses a low-power nanosecond pulsed laser to produce microbubbles. The addition of irradiation from ultrasound causes the microbubbles to expand, collapse, and disrupt the plaque.

"In conventional laser angioplasty, a high laser power is required for the entire cavitation process, whereas in our technology, a lower laser power is only required for initiating the cavitation process," said Singh.

"Overall, the combination of ultrasound and laser reduces the need for laser power and improves the efficiency of atherosclerotic plaque removal."

Because it destroys rather than compresses the plaque, the combination technique will have a lower restenosis rate, or re-narrowing of the artery, compared to balloon angioplasty or stenting. The control provided by the ultrasound and the low-power laser will lower the risk of dissection and perforation in arteries.

The team performed ex vivo experiments on carotid artery plaque samples and pork belly samples, and they are currently planning to

perform in vivo experiments. Both the laser and ultrasound techniques are commonly used by clinicians and should be easy to teach and implement following the in vivo studies.

Combining low-power lasers and ultrasound techniques is not limited to atherosclerosis treatments. Singh and collaborators are also using the methodology for photo-mediated ultrasound therapy and ultrasound-assisted endovascular laser thrombolysis. The former can be used to remove abnormal microvessels in the eye to prevent blindness, while the latter can dissolve blood clots in veins.

Singh will expand upon these additional applications in poster sessions at the ASA meeting, with "Analysis of cavitation induced stresses on blood vessel wall during photo-mediated ultrasound therapy using finite-element based numerical models" and "Combining [ultrasound](#) and endovascular laser for thrombolysis," on May 25, 5-7 p.m. Eastern U.S.

Provided by Acoustical Society of America

Citation: Ultrasound-assisted laser technique vaporizes artery plaque (2022, May 24) retrieved 18 April 2023 from

<https://medicalxpress.com/news/2022-05-ultrasound-assisted-laser-technique-vaporizes-artery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--