

Shorter tuberculosis treatment shows promise in Nigeria

June 13 2022



Credit: Steve Buissinne from Pixabay

Early success with Nigeria's implementation of the new and shorter six-month tuberculosis treatment regimen in four states could pave the way for its rollout nationwide, says an expert.

According to the World Health Organization (WHO), about 10 million people in the world are infected with tuberculosis every year, with

Nigeria and South Africa being among the eight countries that account for two-thirds of all tuberculosis cases.

The previous treatment lasting nine or 18 months was a challenge for patients and [health systems](#) to ensure completion, prompting the WHO and partners to recommend a six-month treatment regimen last month.

Nigeria has already been using the six-month course in four states through an operational research program which began in November 2020.

"So far, our findings reveal that the six-month regimen is cost-effective with mild side effects," says Babawale Victor, assistant director of the National Tuberculosis and Leprosy Control Program at Nigeria's Federal Ministry of Health. "It reduces [drug resistance](#) because of the short period of the treatment, and patients prefer it to the other regimens because recovery is faster."

The six-month regimen either uses BPaLM (a combination of oral medicines bedaquiline, pretomanid, linezolid and moxifloxacin) or BPaL (a combination oral bedaquiline, pretomanid and linezolid).

"With this new guideline, almost all patients with [drug-resistant tuberculosis](#) can be treated in six months with an all-oral regimen," says Tereza Kasaeva, director of the WHO's Global TB Program, announcing the new recommendation last month. "We now have more and much better treatment options for people with [drug resistant] TB thanks to research generating new evidence."

Mel Spigelman, president and CEO of TB Alliance, says that the new WHO guideline allows for the treatment of almost all forms of drug-resistant tuberculosis.

Spigelman tells SciDev.Net: "The WHO has just delivered excellent news for patients. This is a testament to what can be accomplished through long-term, sustained investments in TB research and development.

"We now have a path forward for handling the vast majority of cases of drug-resistant tuberculosis with only six months of all-oral treatment, simplified drug regimens that have manageable side effects, and cure rates on the same order of magnitude as the treatment for drug-sensitive TB."

Spigelman explains that the treatment of drug-resistant tuberculosis could dominate a patient's life and that only a few years ago, treatment for drug-resistant tuberculosis was over three times longer than that of many early-stage cancers.

"With a global treatment success rate of just 59 percent, it was unreliable and treating a patient for almost a year-and-a-half is a long time for them to tolerate the harsh medicines," Spigelman adds.

He says that the new, shorter treatments recommended by the WHO will have an impact on the treatment budget of drug-resistant [tuberculosis](#), and save money.

The new WHO guidelines for drug-resistant TB treatment are expected to be implemented by almost every country in Sub-Saharan Africa, says Spigelman.

He adds that nine countries in Africa including Burkina Faso, Ghana, Malawi, Nigeria and South Africa have made arrangements to procure the medicines directly from manufacturers or the Stop TB Partnership's Global Drug Facility.

Victor tells SciDev.Net of efforts to secure a nationwide rollout of the new treatment.

"At the stakeholders' meeting in June, we will finalize our decision on the rollout of BPaL nationwide," he says. "We will be sending our findings to WHO and await approval of rolling it nationwide."

Provided by SciDev.Net

Citation: Shorter tuberculosis treatment shows promise in Nigeria (2022, June 13) retrieved 19 November 2023 from

<https://medicalxpress.com/news/2022-06-shorter-tuberculosis-treatment-nigeria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.