

By June, almost half of Aussies had recently had COVID-19

July 27 2022



Credit: Pixabay/CC0 Public Domain

It is estimated that by early June, at least 46% of the adult Australian population had acquired infection with SARS-CoV-2, the virus that causes COVID-19, according to results released today from Australia's



most recent serosurvey of antibodies to the virus in blood donors.

The serosurvey, the second in the series of four rounds this year, was conducted by the Kirby Institute at UNSW Sydney and the National Center for Immunization Research and Surveillance (NCIRS) in collaboration with Australian Red Cross Lifeblood, Royal Melbourne Hospital's Victorian Infectious Diseases Reference Laboratory at the Doherty Institute, NSW Health Pathology's Institute of Clinical Pathology and Medical Research (ICPMR), Westmead and other research partners.

The 46.2% prevalence is almost triple what was reported in the previous survey, which estimated that around 17% of the Australian population had been infected by late February.

"While we know there is a lot of virus circulating in the community, we can't rely on testing data alone to understand how many people have been infected, as some may experience only mild symptoms, or none at all, and they may not always have a test to detect the virus. There is also high use of rapid antigen tests, which don't always get reported," says Dr. Dorothy Machalek from the Kirby Institute.

"These serosurveys provide a more complete picture of how much COVID-19 is out there because they measure antibodies to the virus, created when the body's immune system responds to infection, that remain present for many months afterwards."

Evidence of past infection was highest among donors in the 18–29 year age group, at 61.7%, declining with increasing age to 25.7% in donors aged 70–89 years across Victoria, NSW, Queensland and Western Australia. In WA, evidence of past infection in blood donors jumped from 0.5% to 37.5% over a three-month period.



The researchers examined 5,139 de-identified samples from Australian blood donors aged 18–89 years for evidence of COVID-19-related antibodies. Two types of antibody to SARS-CoV-2 were tested for: antibody to the nucleocapsid protein of the virus provides an indication of infection in the past few months, while antibody to the spike protein can indicate past infection and/or vaccination. The <u>blood samples</u> were collected between the 9–18 June 2022.

The researchers noted that the best protection against serious illness from COVID-19 was to be up to date with vaccinations and boosters, in line with current recommendations.

"We are seeing new variants of the omicron version of the virus that causes COVID-19 in circulation in Australia, and that can lead to reinfection even in people who have been infected with an earlier variant" says Professor Kristine Macartney, Director of NCIRS and Professor at the University of Sydney, who is an investigator in the serosurvey program. "Vaccination reduces the risk of severe disease if you get COVID-19, whether it is your first time or a repeat infection."

Professor Paul Kelly, Australia's Chief Medical Officer, says that "data collected from these surveys is crucial for informing our pandemic response."

How are serosurveys conducted?

The ongoing blood <u>donor</u> survey co-led by the Kirby Institute and NCIRS in collaboration with Australian Red Cross Lifeblood also involves investigators at the Victorian Infectious Diseases Reference Laboratory at the Doherty Institute, NSW Health Pathology Institute of Clinical Pathology & Medical Research (ICPMR), the University of Sydney and Murdoch Children's Research Institute.



The residual blood donation samples used in the survey are obtained from Lifeblood's processing centers across the country and are delinked from any identifying information apart from age, sex and post code. Individual results can therefore not be provided back to <u>blood donors</u>.

"Australian Red Cross Lifeblood encourages anyone wanting to contribute to this type of research to become a regular donor. There are many benefits to donating, including finding out your <u>blood</u> type," says Professor David Irving, Director of Research and Development at Australian Red Cross Lifeblood.

The next (third) 2022 round of the Lifeblood donor survey will commence around the end of August. Data is provided to all state and territory governments, and the Commonwealth Government under the Australian National Disease Surveillance Plan for COVID-19.

Provided by University of New South Wales

Citation: By June, almost half of Aussies had recently had COVID-19 (2022, July 27) retrieved 9 July 2023 from <u>https://medicalxpress.com/news/2022-07-june-aussies-covid-.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.