


Researchers reveal the biological basis of network control theory in brain dynamics

December 13 2022




Regional energy efficiency differences between TLE patients and healthy controls. Credit: He Xiaosong et al

A team led by researcher He Xiaosong from the University of Science and Technology (USTC) revealed the correlation between control energy consumption and glucose metabolism in temporal lobe epilepsy (TLE), providing the biological basis for the application of network control theory (NCT) in the study of brain dynamics. This work was published in *Science Advances* on November 9.

Understanding the mechanism of neural dynamics in various different [brain regions](#) has been a [major focus](#) in neuroscience. In recent years, NCT, a theory that originates in engineering, has been widely used to estimate the control energy in the dynamic processes in the brain by modeling the transitions of different brain states. The control energy originally referred to the energy input required to change the state of the system. However, when it comes to brain dynamics, the biological counterpart of such energy input remains unknown, which raises questions about the application of NCT.

Based on a comparison between the brain structures of TLE patients and healthy controls, the team simulated two representative brain dynamic processes and calculated the control energy consumed in these processes. The results showed that patients with TLE required significantly higher global optimal control energy (OCE) than healthy controls to activate the limbic network compared to healthy controls. The abnormal energetic efficiency is consistent with the lateralization of the seizure focus, which implies that the seven limbic brain regions ipsilateral to the seizure focus consume more control energy to maintain brain dynamics.



Correlation between baseline glucose metabolism and control energy consumption. Credit: HE Xiaosong et al.

Using [positron emission tomography](#), the team confirmed that the baseline glucose metabolism levels in these brain regions are inversely correlated with the consumption level of control energy, which means that lower baseline metabolism requires more energy to obtain the same

level of activation. A decline in baseline regional metabolism arose from the loss of structural integrity, especially in the hippocampus, and eventually led to higher energy costs for maintaining brain dynamics.

The team delivered a unified framework for the correlations between the structural integrity of brains, the baseline [glucose metabolism](#) and the control energy required to maintain brain dynamics. By providing a potential biological explanation of control energy, this work lays the foundation for further NCT applications in neuroscience.

More information: Xiaosong He et al, Uncovering the biological basis of control energy: Structural and metabolic correlates of energy inefficiency in temporal lobe epilepsy, *Science Advances* (2022). [DOI: 10.1126/sciadv.abn2293](https://doi.org/10.1126/sciadv.abn2293)

Provided by University of Science and Technology of China

Citation: Researchers reveal the biological basis of network control theory in brain dynamics (2022, December 13) retrieved 3 July 2023 from <https://medicalxpress.com/news/2022-12-reveal-biological-basis-network-theory.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--