


Could antidepressants cause treatment-emergent mania?

February 27 2023, by Colette Gallagher



Increased rates of TEM+ with antidepressants that increase mitochondrial function. Rates of treatment-induced mania with antidepressants that increase (Mito+) were two times more frequent than with those that decrease (Mito-) mitochondrial function [TEM+ Mito+ = 24.7%; TEM+ Mito- = 13.5%; OR = 2.21; $p = 0.000009$]. Credit: *Molecular Psychiatry* (2022). DOI: 10.1038/s41380-022-01888-x

In a study published in *Molecular Psychiatry*, Mark Frye, M.D., a Mayo Clinic researcher and collaborators, investigated the risk of treatment-

emergent mania in bipolar disorder when treated with antidepressants.

"We found that [antidepressants](#) that increase mitochondrial energetics (cells that extract energy from nutrients for sustaining life) may elevate the risk of treatment-emergent mania," says Dr. Frye.

The increased energy expenditure of mania associated with impulsivity, poor judgment, psychosis and loss of insight can drive high-risk behaviors, often resulting in hospitalization or incarceration. The aftermath of mania can have an enduring negative effect on the patient's quality of life, explains Dr. Frye, the study's senior author.

Dr. Frye specializes in the neurobiology of bipolar disorders. Bipolar disorder, formerly called [manic depression](#), is a [mental health condition](#) that causes [extreme mood swings](#) that include emotional highs (mania or hypomania) and lows (depression).

"These data suggest categorizing antidepressants based on mitochondrial energetics may be of value," says Dr. Frye.

The clinical practice is of significant consequence, as antidepressant prescriptions for bipolar disorder in the U.S. have more than doubled in the last two decades from 17.9% to 40.9%, expounds Dr. Frye.

The study assessed 692 participants 18 to 80 years old from the Mayo Clinic Bipolar Disorder Biobank with a known history of antidepressant exposure. The researchers found that compared to participants exposed to antidepressants that decrease mitochondrial activity, treatment-emergent mania was twice as common in patients exposed to antidepressants that increase mitochondrial energetics.

"Our study provides early evidence that supports the hypothesis of an amplified response in mitochondrial energetics of select antidepressants

that drive, in part, the cause of treatment-emergent mania," says Dr. Frye. "These data suggest that categorizing antidepressants based on mitochondrial energetics warrants further large-scale clinical and pharmacogenomic studies to determine the effect on managing and treating patients with [bipolar disorder](#)."

More information: Manuel Gardea-Resendez et al, Antidepressants that increase mitochondrial energetics may elevate risk of treatment-emergent mania, *Molecular Psychiatry* (2022). [DOI: 10.1038/s41380-022-01888-x](#)

Provided by Mayo Clinic

Citation: Could antidepressants cause treatment-emergent mania? (2023, February 27) retrieved 8 April 2023 from <https://medicalxpress.com/news/2023-02-antidepressants-treatment-emergent-mania.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--