


# Machine learning identifies 'heart roundness' as a new tool for diagnosing cardiovascular conditions

March 29 2023



Deep learning-enabled analysis of medical images identifies cardiac sphericity as an early marker of cardiomyopathy and related outcomes. Credit: Med/Vukadinovic et al.

Physicians currently use assessments like heart chamber size and systolic function to diagnose and monitor cardiomyopathy and other related heart conditions. A paper published in the journal *Med* on March 29 suggests that another measurement—cardiac sphericity, or roundness of the heart—may one day be a useful implement to add to the diagnostic toolkit.

"Roundness of the [heart](#) isn't necessarily the problem per se—it's a marker of the problem," says co-corresponding author Shoa L. Clarke, a preventive cardiologist and an instructor at Stanford University School of Medicine. "People with rounder hearts may have underlying cardiomyopathy or underlying dysfunction with the molecular and cellular functions of the heart muscle. It could be reasonable to ask whether there is any utility in incorporating measurements of sphericity into clinical decision-making."

This proof-of-concept study used [big data](#) and machine learning to look at whether other anatomical changes in the heart could improve the understanding of cardiovascular risk and pathophysiology. The investigators chose to focus on sphericity because [clinical experience](#) had suggested it is associated with [heart problems](#). Prior research had primarily focused on sphericity after the onset of heart disease, and they hypothesized that sphericity may increase even before the onset of clinical heart disease.

"We have established traditional ways of evaluating the heart, which have been important for how we diagnose and treat [heart disease](#)," Clarke says. "Now with the ability to use deep-learning techniques to look at [medical images](#) at scale, we have the opportunity to identify new ways of evaluating the heart that maybe we haven't considered much in the past."

"They say a picture is worth a thousand words, and we show that this is very true for [medical imaging](#)," says co-corresponding author David Ouyang, a cardiologist and researcher at the Smidt Heart Institute of Cedars-Sinai. "There's a lot more information available than what physicians are currently using. And just as we've previously known that a bigger heart isn't always better, we're learning that a rounder heart is also not better."

This research employed data from the UK Biobank, which includes genetic and clinical information on 500,000 people. As part of that study, a subset of volunteers had MRI imaging of their hearts performed. The California-based team used data from a subset of about 38,000 UK Biobank study participants who had MRIs that were considered normal at the time of the scans. Subsequent medical records from the volunteers indicated which of them later went on to develop diseases like cardiomyopathy, atrial fibrillation, or heart failure and which did not.

The researchers then used deep-learning techniques to automate the measurement of sphericity. Increased cardiac sphericity appeared to be linked to future heart troubles.

The investigators also looked at genetic drivers for cardiac sphericity and found overlap with the genetic drivers for cardiomyopathy. Using Mendelian randomization, they were able to infer that intrinsic disease of the heart muscle—meaning defects not caused by heart attacks—caused cardiac sphericity.

"There are two ways that these findings could add value," Ouyang says. "First, they might allow physicians to gain greater clinical intuition on how patients are likely to do at a very rapid glance. In the broader picture, this research suggests there are probably many useful measurements that clinicians still don't understand or haven't discovered. We hope to identify other ways to use imaging to help us predict what will happen next."

The researchers emphasize that much more research is needed before the findings from this study can be translated to clinical practice. For one thing, the connection is still speculative and would need to be confirmed with additional data. If the link is confirmed, a threshold would need to be established to indicate what degree of sphericity might suggest that clinical interventions are needed. The team is sharing all the data from this work and making them available to other investigators to begin answering some of these questions.

Additionally, ultrasound is more commonly used than MRI to image the heart. To further advance this research, replicating these findings using ultrasound images will be useful, they note.

**More information:** Shoa L. Clarke, Deep learning-enabled analysis of medical images identifies cardiac sphericity as an early marker of cardiomyopathy and related outcomes, *Med* (2023). [DOI: 10.1016/j.medj.2023.02.009](https://doi.org/10.1016/j.medj.2023.02.009).  
[www.cell.com/med/fulltext/S2666-6340\(23\)00069-7](https://www.cell.com/med/fulltext/S2666-6340(23)00069-7)

Provided by Cell Press

Citation: Machine learning identifies 'heart roundness' as a new tool for diagnosing cardiovascular conditions (2023, March 29) retrieved 20 November 2023 from

<https://medicalxpress.com/news/2023-03-machine-heart-roundness-tool-cardiovascular.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.