


Study uncovers novel synaptic intricacies inside the retina

June 19 2023, by Melissa Rohman



Array of cone photoreceptor terminals labeled for PSD95 (red), the glutamate receptors belonging to one type of bipolar cell (yellow), and glutamate re-uptake transporters (cyan). Credit: Steven DeVries, MD, PhD

A Northwestern Medicine study has uncovered novel cellular mechanisms within the retina, findings that could help advance the development of targeted therapeutics for diseases and conditions impacting vision, according to findings published in *Nature Communications*.

Cone synapses inside the eye's retina help the brain process changes in light. It's a unique synapse because it has evolved to signal changes in [light intensity](#), said Steven DeVries, MD, Ph.D., the David Shoch, MD, Ph.D., Professor of Ophthalmology and senior author of the study.

"Counterintuitively, cone neurotransmitter release is high in the dark and reduced by light. When the light's brighter, the reduction is larger. When the lights are dimmer, it's smaller; it operates differently from most synapses which use an increase in transmitter release to signal all-or-nothing, digital action potentials," DeVries said.

Unlike most other synapses in the brain, each individual cone synapse is connected to more than a dozen different types of post-synaptic neurons, the [bipolar cells](#), which relay information in parallel to the inner retina. In the inner retina, these parallel streams not only contribute to conscious vision but also to subconscious processes like gaze stabilization.

In the current study, which involved non-human mammalian retinas, the investigators first used super-resolution microscopy to map out the locations of transmitter release sites, transmitter re-uptake proteins and

post-synaptic contacts at the cone synapse. Then they used an approach called "synaptic accounting," to relate the amount of transmitter released by a cone to the responses in each post-synaptic bipolar cell type.

"Transmitter is released in packets or quanta when a vesicle fuses with the presynaptic membrane. Since most synapses involve one-to-one direct contacts across a narrow cleft, it is assumed that one detected quantum equals one released quantum. The cone synapse has a different design that voids this assumption. We developed a way to stimulate a cone and count the vesicles that are released while at the same time counting the number of vesicles that are detected by the postsynaptic neuron," DeVries said.

Using these techniques, the investigators showed that certain bipolar cell types respond to individual fusion events and total quanta while other types respond to degrees of locally coincident events, creating a nonlinear summation. These differences are caused by a combination of factors specific to each bipolar cell type, including diffusion distance, contact number, receptor affinity, and proximity to transporters.

"The outer [retina](#) uses the same toolbox as elsewhere in the [central nervous system](#), like vesicles, synaptic release zones and postsynaptic receptors, but organizes these elements in novel ways to accomplish a different, very localized type of processing. Analog processing is also found in the dendritic tree of central nervous system neurons, where the bulk of calculation, both linear and nonlinear, occurs," DeVries said.

According to DeVries, one next step for his team includes using a new, more powerful type of [super-resolution microscopy](#) to determine the protein components that make up cone synapses.

"One of the ways that the different bipolar cells divide the cone signal up is that some of them are very sensitive to small signals and others require

strong signals to respond; the 'strong signal' or high threshold bipolar cell has a unique type of insensitive post-synaptic receptor. We would also like to identify this receptor," DeVries said.

More information: Chad P. Grabner et al, Mechanisms of simultaneous linear and nonlinear computations at the mammalian cone photoreceptor synapse, *Nature Communications* (2023). [DOI: 10.1038/s41467-023-38943-2](https://doi.org/10.1038/s41467-023-38943-2)

Provided by Northwestern University

Citation: Study uncovers novel synaptic intricacies inside the retina (2023, June 19) retrieved 4 July 2023 from <https://medicalxpress.com/news/2023-06-uncovers-synaptic-intricacies-retina.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.