

Second year of COVID pandemic brought spike in child mental health visits to ED

July 13 2023



Credit: Unsplash/CC0 Public Domain

The number of young people in the United States visiting hospital emergency departments for mental health crises increased sharply during the second year of the COVID-19 pandemic, according to a study led by researchers from the Department of Health Care Policy in the Blavatnik Institute at Harvard Medical School. The findings are published Jul 12 in *JAMA Psychiatry*.

These results come amidst growing national concern about a crisis in youth mental [health](#) and provide important new details about how [young people](#) with [mental health problems](#) such as [self-harm](#) and suicide attempts are using acute medical services.

The findings, the researchers said, highlight the critical need for policies to increase resources for mental health for all aspects of care, including emergency departments, inpatient pediatric mental health facilities, [primary care](#), and prevention.

"The bottom line is that as a society, we need to do more to protect the mental health and well-being of our young people," said Haiden Huskamp, Henry J. Kaiser Professor of Health Care Policy at HMS.

Numerous reports have noted that the stress and isolation of the COVID-19 [pandemic](#) have exacerbated what US Surgeon General Vivek Murthy has [described as a crisis](#) of adolescent mental health.

And the trend is not new, as numerous studies have shown. The suicide rate among young people [increased by 57 percent](#) in the decade before the pandemic, compared with the preceding decade. With increasing prevalence of mental illness among youth and a chronic lack of providers, the mental health care system has been stressed for a long time, the researchers said.

The pandemic helped bring those festering problems to a head, the authors said. The multiple and compounding stressors of COVID-19 have taken a grave toll on the mental health of a whole generation of young people and are taxing a mental health care system that's already stretched to capacity, they said.

"One of the most concerning findings was the dramatic increase in the number of adolescents waiting multiple days in the emergency room

before being admitted to facilities that can provide the level of treatment they need," said Huskamp.

For their analysis, the researchers looked at private health insurance claims submitted between March 2019 and February 2022 for more than 4 million people between the ages of 5 and 17. The researchers compared numbers and outcomes of emergency department visits related to mental health conditions from the year before the COVID-19 pandemic (March 2019 to February 2020) with data from the second year of the pandemic (March 2021 to February 2022).

The young people in the study sample were 7 percent more likely to have had an ED visit for mental health in the second year of the pandemic than they were in the 12 months prior to the pandemic. The overall increase was driven by a dramatic surge in emergency department visits among [adolescent girls](#), who were 22 percent more likely to have an [emergency room](#) visit during the second year of the pandemic compared with the year before the virus hit.

"One surprising and concerning finding was that the increase in ED visits was largely driven by girls who came to the hospital for conditions such as suicidal thoughts or plans, [suicide attempts](#), and self-harm," first author Lindsay Overhage, an HMS MD/Ph.D. student with an interest in mental health policy, said. "It's critical that we do all we can to prevent these serious illnesses and to treat those who are suffering."

Overall, the likelihood that a child who visited the ED for mental health care would be admitted to an inpatient mental health program increased by 8 percent in the second year of the pandemic, relative to the year before the outbreak. The number of young people who spent at least two days waiting to be admitted from the ED to an inpatient psychiatric service increased by 76 percent.

The findings underscore an urgent need to identify and relieve the underlying stresses that are driving this steep rise in depression, anxiety, self-harm, and other serious mental health problems among young people in an effort to prevent suffering, the researchers said. These efforts, they added, must include research to help understand why girls have been affected worse than boys.

The study also highlights the importance of working rapidly to increase inpatient and outpatient child psychiatry capacity to give young people in crisis the care that they need and to reduce the strain on the acute mental health care system, the researchers said. The researchers point to a variety of ways to address this problem including improving inpatient capacity, increasing the availability of mental health providers, preventing and fighting burnout among mental health care providers, and supporting non-specialist primary care and emergency care clinicians who provide [mental health](#) care.

For children in crisis now, the researchers note that there are promising treatments that can be delivered in emergency departments, in person or using telemedicine. These therapies may reduce the need for hospital admissions or at least allow patients to begin some effective treatment while they are waiting for a spot in an inpatient program.

More information: Lindsay Overhage et al, Trends in Acute Care Use for Mental Health Conditions Among Youth During the COVID-19 Pandemic, *JAMA Psychiatry* (2023). [DOI: 10.1001/jamapsychiatry.2023.2195](#)

Provided by Harvard Medical School

Citation: Second year of COVID pandemic brought spike in child mental health visits to ED

(2023, July 13) retrieved 13 July 2023 from <https://medicalxpress.com/news/2023-07-year-covid-pandemic-brought-spike.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.