

PET scan distinguishes Alzheimer's from other dementia

1 November 2007

A PET scan (positron emission tomography) that measures uptake of sugar in the brain significantly improves the accuracy of diagnosing a type of dementia often mistaken for Alzheimer's disease, a study led by a University of Utah dementia expert has found.

The scan, FDG-PET, helped six doctors from three national Alzheimer's disease centers correctly diagnose frontotemporal dementia (FTD) and Alzheimer's in almost 90 percent of cases in the study—an improvement of as much as 14 percent from usual clinical diagnostic methods. FDG stands for fluorodeoxyglucose, a short-lived radioactive form of sugar injected into people during PET scans to show activity levels in different parts of the brain. In Alzheimer's, low activity is mostly in the back part of the brain; in FTD, low activity is mostly in the front of the brain.

FDG-PET is an especially powerful tool in early treatment of FTD, said the study's lead author, Norman L. Foster, M.D., professor of neurology and director of the Center for Alzheimer's Care, Imaging and Research at the University of Utah School of Medicine.

FTD is a common cause of early onset dementia among people 45-64 years old and is marked by behavioral changes and language difficulties. Like Alzheimer's, it can take years to develop and, for now, is incurable. Although FTD is a separate disorder, it often meets clinical diagnostic criteria for Alzheimer's and often is misdiagnosed even by dementia experts.

"Early diagnosis of FTD can have a tremendous impact on the treatment for patients and their family members. Many patients are misdiagnosed and may be hospitalized and receive drugs for the wrong disease," Foster said. "Accurate diagnosis bypasses the costs, side-effects, and frustration of misguided care. Furthermore, one-third of FTD patients have a family history of a similar disorder

and family members need to know if they are at increased risk of the disease."

The study was funded by the National Institute on Aging, a part of the National Institutes of Health.

"Dr. Foster's work involving patients from several NIA-sponsored Alzheimer's Disease Centers advances the use of PET imaging as a clinical tool," said Creighton Phelps, Ph.D., program director of the Alzheimer Disease Centers at the National Institute on Aging. "Combined with the patient's medical history and psychometric testing, it enhances a physician's ability to more accurately distinguish between FTD and early-onset AD."

As the U.S. population ages, the number of people with dementia is projected to increase markedly, with Utah and the Intermountain West expected to experience the fastest rate of growth. Although FDG-PET is widely available, it is not often used in dementia, because of insurance concerns. Medicare recently agreed to pay for FDG-PET scans to evaluate dementia, but currently many insurance companies in the Intermountain West and Utah do not. Foster is working to make FDG-PET scans available to those who need them and results of this study prove they sometimes are worth doing.

"This study shows FDG-PET is a reliable and valid diagnostic test that can aid physicians in making the sometimes difficult clinical distinction between AD (Alzheimer's disease) and FTD," Foster and his co-authors wrote. But PET scans alone are not enough to confirm FTD or Alzheimer's. "A careful consideration of the medical history and examination will continue to be essential to dementia evaluation."

The study has appeared online in the journal *Brain*.

Foster and his colleagues examined the medical records and FDG-PET scans of 45 patients who

later had autopsies. Microscopic examination found 31 had Alzheimer's and 14 had FTD. The researchers summarized the clinical course of the disease in each patient. The expert neurologists at the NIH centers, who had 10 years to 25 years of experience, then were asked to decide what caused each patient's dementia using clinical information alone or using FDG-PET images.

The experts correctly distinguished FTD and Alzheimer's using only the clinical methods in 76 percent to 79 percent of the cases. Using the FDG-PET scans alone, however, the physicians correctly diagnosed the two dementias in 85 percent to 89 percent of cases. Adding FDG-PET to clinical information increased the correct diagnosis from 79 percent to 90 percent. The highest accuracy in diagnosis was achieved with SSP (stereotactic surface projection) displays, which summarize changes in brain activity and apply a statistical test to show significant areas of damage.

The PET scans also had other benefits. The researchers found in 42 percent of cases, the scans increased the experts' confidence in a correct diagnosis or made them question and sometimes change an incorrect diagnosis.

Source: University of Utah

APA citation: PET scan distinguishes Alzheimer's from other dementia (2007, November 1) retrieved 1 July 2022 from

<https://medicalxpress.com/news/2007-11-pet-scan-distinguishes-alzheimer-dementia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.