

IL-22 gene delivers the goods and decreases intestinal inflammation

2 January 2008

There are two major types of inflammatory bowel disease (IBD), Crohn disease (CD) and ulcerative colitis (UC). Conflicting reports have indicated that the soluble factor IL-22 can have both IBD promoting and IBD controlling effects. But now, Atsushi Mizoguchi and colleagues at Massachusetts General Hospital, Boston, have established that IL-22 ameliorates disease in a mouse model of UC.

Expression of IL-22 is much higher in the intestines of individuals with CD than UC. To investigate the role of IL-22 in IBD, the authors used a new microinjection-based strategy to deliver the gene that makes IL-22 to the walls of the intestine of mice who suffer from an intestinal disease that models UC. Delivery of the IL-22 gene ameliorated local intestinal inflammation through enhanced mucus production. Consistent with this, when the same strategy was used to deliver a gene that makes a protein that neutralizes IL-22, IL-22-binding protein, to the walls of the intestines of normal mice it enhanced chemical-induced intestinal inflammation. The authors therefore suggest that individuals with UC might benefit from local delivery of the IL-22 gene to their intestines.

Source: Journal of Clinical Investigation

APA citation: IL-22 gene delivers the goods and decreases intestinal inflammation (2008, January 2) retrieved 18 June 2022 from <https://medicalxpress.com/news/2008-01-il-gene-goods-decreases-intestinal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.