

Study examines prevalence of chest pain in patients 1 year after heart attack

23 June 2008

Nearly one in five patients experiences chest pain one year after having a heart attack, according to a report in the June 23 issue of *Archives of Internal Medicine*, one of the JAMA/Archives journals.

continue smoking, to undergo revascularization (surgery to reestablish blood flow to the heart) after hospitalization and to have significant new, persistent or fleeting depressive symptoms.

One of the main goals of in-hospital treatment and outpatient care after heart attack is to relieve angina (or episodic chest pain), according to background information in the article. The prevalence and treatment of chest pain one year after heart attack are largely unknown. "By identifying these factors, a more complete understanding of those patients who are at the greatest risk for angina [chest pain] after myocardial infarction [heart attack] can occur," the authors write. Identifying this population is important for treating remaining chest pain and improving patient outcomes, including ability to exercise and health-related quality of life.

Source: JAMA and Archives Journals

Thomas M. Maddox, M.D., S.M., of Denver Veterans Affairs Medical Center and University of Colorado Denver, and colleagues studied the occurrence of angina in 1,957 patients recruited from January 2003 to June 2004. Patients filled out questionnaires assessing their chest pain one year after hospitalization for heart attack. Sociodemographic, clinical and other lifestyle factors were also reported.

Of all patients, 389 (19.9 percent) reported angina one year after hospitalization for heart attack.

Twenty-four patients (1.2 percent) reported having daily chest pain, 59 (3 percent) reported weekly chest pain and 306 (15.6 percent) reported having chest pain less than once a week.

Patients experiencing chest pain one year after heart attack were more likely to be younger, non-white males with prior chest pain who have undergone prior coronary artery bypass graft surgery and have experienced recurring rest chest pain while hospitalized for heart attack. Patients with one-year chest pain were also more likely to

APA citation: Study examines prevalence of chest pain in patients 1 year after heart attack (2008, June 23) retrieved 5 May 2021 from <https://medicalxpress.com/news/2008-06-prevalence-chest-pain-patients-year.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.