

Study finds international variations in quality of life loss after fracture

24 March 2011

A study presented today at the European Congress on Osteoporosis and Osteoarthritis currently taking place in Valencia, Spain has found that the initial quality of life loss following an osteoporotic fracture is substantial, especially with regard to hip and vertebral fractures. The study found differences in quality of life loss between countries after correcting for other explanatory variables.

[DOI 10.1007/s00198-011-1554-9](https://doi.org/10.1007/s00198-011-1554-9)

Provided by International Osteoporosis Foundation

The [quality of life](#) (QoL) of patients who sustained a hip, wrist or vertebral fracture was examined for the four months following the fracture. The study is part of the International Costs and Utilities Related to Osteoporotic Fractures Study (ICUROS) which is an international patient-based study endorsed by the International Osteoporosis Foundation. ICUROS investigates the costs and [health effects](#) of osteoporotic fractures in women and men over the age of fifty. The aim is to gather better information regarding the burden of [osteoporosis](#) in the participating countries and worldwide. Currently, eleven countries are involved in ICUROS.

Involving 2,737 patients from nine countries, the study found that quality of life loss over four months was highest for hip fractures, followed by vertebral and wrist fractures. Interestingly, significant variations between countries were identified. Quality of life loss after hip fracture was greatest in Lithuania, followed by Italy, Russia and Mexico. Lower quality of life loss was experienced by hip fracture patients in Austria and Sweden. In all countries in which vertebral fractures were studied, patients who were hospitalized experienced more quality of life loss than those who were not hospitalized following vertebral fracture.

More information: The abstract 'Quality of life four months following a fracture - results from the ICUROS', F. Borgström, et al. (OC14) is published in supplement issue of Osteoporosis International.

APA citation: Study finds international variations in quality of life loss after fracture (2011, March 24) retrieved 9 July 2022 from <https://medicalxpress.com/news/2011-03-international-variations-quality-life-loss.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.