

Low vitamin D levels do not increase the risk of chronic obstructive pulmonary disease

4 November 2011

Vitamin D levels are not related to acute exacerbations of chronic obstructive pulmonary disease (AECOPD) in patients with severe COPD, according to a large prospective cohort study involving 973 North American patients. The findings were published online ahead of the print edition of the *American Thoracic Society's American Journal of Respiratory and Critical Care Medicine*.

In the study, a secondary analysis of data from a randomized, controlled trial of the effects of azithromycin on the frequency of AECOPD, no relationship was found between baseline [Vitamin D](#) levels and time to first AECOPD or between vitamin D levels and AECOPD exacerbation rates.

"Vitamin D insufficiency and deficiency are common in patients with COPD, and patients with severe COPD are at the highest risk for exacerbations, so we hypothesized that low vitamin D levels might increase the risk of AECOPDs," said Ken M. Kunisaki, M.D., of the Minneapolis Veterans Affairs Medical Center. "Our negative results are in contrast with earlier studies in which lower vitamin D levels were associated with higher rates of respiratory infections in adults and more frequent [asthma exacerbations](#) in children."

In the current study, mean forced expiratory volume in one second ([FEV1](#)) was 1.12L, 40% of predicted. Mean vitamin D level at baseline was 25.7 ± 12.8 ng/mL, with 33.1% of subjects categorized as vitamin D insufficient (<20 ng/mL but

APA citation: Low vitamin D levels do not increase the risk of chronic obstructive pulmonary disease (2011, November 4) retrieved 10 November 2022 from <https://medicalxpress.com/news/2011-11-vitamin-d-chronic-obstructive-pulmonary.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.