

# Modest alcohol intake associated with less inflammation in patients with common liver disease

May 15 2012

---

NAFLD (non-alcoholic fatty liver disease) is the most common type of liver disease in the developed world, affecting up to one-third of the US population. NAFLD is often associated with obesity and other parameters of the so-called "metabolic syndrome," which is a major risk factor for the development of cardiovascular disease. In a well-done study among subjects with NAFLD, the investigators have demonstrated that modest alcohol consumption (an average of up to 20 grams of alcohol per day and no binge drinking) is associated with less evidence of inflammation of the liver (steatohepatitis), a condition known to markedly increase the risk of progression of liver disease to cirrhosis.

Given that NAFLD and other conditions associated with the metabolic syndrome are so common, and are major risk factors for developing cardiovascular disease, the results of the present study are important. They show that modest drinking is associated with decreased, not increased, inflammation of the liver. Further, even among subjects with NAFLD, cardiovascular disease is a much more common cause of death than liver disease. The authors suggest that intervention studies should be done to support their findings; if confirmed, subjects with NAFLD should not be advised to avoid all alcohol, which is the current advice usually given to such patients.

**More information:** Dunn W, Sanyal AJ, Brunt EM, Unalp-Arida A, Donohue M, McCullough AJ, Schwimmer JB. Modest alcohol

consumption is associated with decreased prevalence of steatohepatitis in patients with non-alcoholic fatty liver disease (NAFLD). *Journal of Hepatology* 2012 (pre-publication release)

Provided by Boston University Medical Center

Citation: Modest alcohol intake associated with less inflammation in patients with common liver disease (2012, May 15) retrieved 21 December 2023 from <https://medicalxpress.com/news/2012-05-modest-alcohol-intake-inflammation-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.