


# Oxytocin improves brain function in children with autism

21 May 2012, By Karen N. Peart



(Medical Xpress) -- Preliminary results from an ongoing, large-scale study by Yale School of Medicine researchers shows that oxytocin - a naturally occurring substance produced in the brain and throughout the body- increased brain function in regions that are known to process social information in children and adolescents with autism spectrum disorders (ASD).

A Yale Child Study Center research team that includes postdoctoral fellow Ilanit Gordon and Kevin Pelphrey, the Harris Associate Professor of Child Psychiatry and Psychology, presented the results on Saturday, May 19 at the International Meeting for Autism Research.

"Our findings provide the first, critical steps toward devising more effective treatments for the core social deficits in autism, which may involve a combination of clinical interventions with an administration of [oxytocin](#)," said Gordon. "Such a treatment approach will fundamentally improve our understanding of autism and its treatment."

Social-communicative dysfunctions are a core characteristic of autism, a neurodevelopmental disorder that can have an enormous emotional and financial burden on the affected individual, their

families, and society.

Gordon said that while a great deal of progress has been made in the field of [autism](#) research, there remain few effective treatments and none that directly target the core social dysfunction. Oxytocin has recently received attention for its involvement in regulating social abilities because of its role in many aspects of social behavior and social cognition in humans and other species.

To assess the impact of oxytocin on the [brain function](#), Gordon and her team conducted a first-of-its-kind, double-blind, placebo-controlled study on children and adolescents aged 7 to 18 with ASD. The team members gave the children a single dose of oxytocin in a nasal spray and used functional magnetic resonance brain imaging to observe its effect.

The team found that oxytocin increased activations in brain regions known to process social information. Gordon said these brain activations were linked to tasks involving multiple social information processing routes, such as seeing, hearing, and processing information relevant to understanding other people.

Provided by Yale University

APA citation: Oxytocin improves brain function in children with autism (2012, May 21) retrieved 8 October 2022 from <https://medicalxpress.com/news/2012-05-oxytocin-brain-function-children-autism.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*