

Revisiting the association between saturated fat intake and coronary heart disease

July 3 2012

In this week's *PLoS Medicine*, Kay-Tee Khaw of the University of Cambridge, UK and colleagues analyze data from a prospective cohort study and show associations between plasma concentrations of saturated phospholipid fatty acids and risk of coronary heart disease, and an inverse association between omega-6 polyunsaturated phospholipid fatty acids and risk of coronary heart disease.

The authors comment: "Early guidelines to prevent [coronary heart disease] recommended reductions in saturated fat but little consistency as to what might be substituted: other fats, protein, or carbohydrate. Our results add to the accumulating evidence that substitution of saturated fat by n-6 polyunsaturated fat may have more [[coronary heart disease](#)] benefits."

More information: Khaw K-T, Friesen MD, Riboli E, Luben R, Wareham N (2012) Plasma Phospholipid Fatty Acid Concentration and Incident Coronary Heart Disease in Men and Women: The EPIC-Norfolk Prospective Study. *PLoS Med* 9(7): e1001255.
doi:10.1371/journal.pmed.1001255

Provided by Public Library of Science

Citation: Revisiting the association between saturated fat intake and coronary heart disease (2012, July 3) retrieved 13 January 2023 from

<https://medicalxpress.com/news/2012-07-revisiting-association-saturated-fat-intake.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.