


Passive smoking increases platelet activation in healthy people

27 August 2012



The [blood levels](#) of all three parameters were statistically higher after subjects were exposed to passive smoking (see figure 1). COHb increased from $0.8 \pm 0.3\%$ at baseline to $1.2 \pm 0.4\%$ after one hour in a smoking room ($p=0.001$). Lactate increased from 0.70 ± 0.2 mmol/L to 2.2 ± 0.9 mmol/L ($p=0.001$). MPV increased from 7.8 ± 0.4 femtolitres (fL) to 8.5 ± 0.6 fL ($p=0.001$). Dr Kaya said: "These results show that passive smoking increases platelet activation and increases CO and lactate levels in the blood."

The researchers found significant correlations between MPV and COHb levels ($r=0.55$, $p=0.0001$) and between MPV and lactate levels ($r=0.65$, $p=0.0001$) after smoking. There was also a remarkable relation between COHb and lactate levels after smoking ($r=0.78$, $p=0.0001$). Dr Kaya said: "Previous studies have suggested that the chemicals in cigarette smoke, especially nicotine and CO, increase platelet-activating factor. The correlations found in our study suggest that the CO in cigarette smoke also increases MPV levels. It is likely that [lactate](#) levels increased because oxygen levels in the blood dropped as CO increased."

He concluded: "We have shown that 1 hour exposure to passive smoking increases platelet activation, which could be the mechanism by which it contributes to increased risk of thrombotic events in healthy people. It is likely that prolonged exposure to passive smoking could have even greater effects. Healthy people should avoid exposure to passive smoking so that they do not increase their risk of thrombotic events."

"It is well known that passive smoking is harmful for cardiovascular health, but the mechanism has not yet been discovered," said Dr Kaya. "We investigated the effects of passive smoking on the levels of three parameters – mean platelet volume (MPV), carboxyhemoglobin (COHb) and lactate - in an effort to further understand this mechanism. We also looked at the correlation between the three parameters."

Mean platelet volume (MPV) is a well established indicator of platelet activation that is increased in acute thrombotic events. It is suspected that carbon monoxide (CO) plays a major role in cigarette smoke-induced cardiovascular diseases. When CO binds with [hemoglobin](#) (Hb) in [red blood cells](#) it is called COHb. Lactate accumulates in the blood when the supply of oxygen to blood cells is limited.

For the study, 55 healthy nonsmoker volunteers (males 30%; mean age 26 ± 5 years) were prospectively enrolled. [Blood samples](#) for measurements of MPV, COHb and lactate were taken at baseline and after spending one hour in a [smoking room](#).

Provided by European Society of Cardiology

APA citation: Passive smoking increases platelet activation in healthy people (2012, August 27) retrieved 1 June 2022 from <https://medicalxpress.com/news/2012-08-passive-platelet-healthy-people.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.