

Clopidogrel after MI less effective in diabetes patients

5 September 2012



Clopidogrel therapy following a heart attack does less to reduce the risk of death in patients with diabetes than in those without diabetes, according to a study published in the Sept. 5 issue of the *Journal of the American Medical Association*.

(HealthDay)—Clopidogrel therapy following a heart attack does less to reduce the risk of death in patients with diabetes than in those without diabetes, according to a study published in the Sept. 5 issue of the *Journal of the American Medical Association*.

Charlotte Andersson, M.D., Ph.D., from Gentofte Hospital in Hellerup, Denmark, and colleagues analyzed data from the Danish nationwide administrative registries (2002 to 2009) for 58,851 patients hospitalized with incident [myocardial infarction](#) (MI) who had survived and had not undergone [coronary artery bypass surgery](#) 30 days after discharge.

The researchers found that recurrent MI occurred in 1,790 patients with diabetes (25 percent) and 7,931 patients without diabetes (15 percent); of these, 1,225 with (17 percent) and 5,377 without diabetes (10 percent) died. The unadjusted mortality rates (events/100 person-years) were 13.4 (95 percent confidence interval [CI], 12.8 to 14.0) versus 29.3 (95 percent CI, 28.3 to 30.4) for patients with diabetes treated with clopidogrel versus those not treated. The unadjusted mortality rates were 6.4 (95 percent CI, 6.3 to 6.6) versus

21.3 (95 percent CI, 21.0 to 21.7) for those patients without diabetes treated with clopidogrel versus those not treated. Clopidogrel was associated with significantly less effectiveness for all-cause mortality (hazard ratio, 0.89 [95 percent CI, 0.79 to 1.00] versus 0.75 [95 percent CI, 0.70 to 0.80]) and for [cardiovascular mortality](#) (hazard ratio, 0.93 [95 percent CI, 0.81 to 1.06] versus 0.77 [95 percent CI, 0.72 to 0.83]) among patients with diabetes versus those without diabetes.

"Among patients with diabetes compared with patients without diabetes, the use of conventional clopidogrel treatment after MI was associated with lower reduction in the risk of all-cause death and cardiovascular death," the authors write.

Several authors disclosed financial ties to the pharmaceutical industry.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)
[Editorial \(subscription or payment may be required\)](#)

Copyright © 2012 [HealthDay](#). All rights reserved.

APA citation: Clopidogrel after MI less effective in diabetes patients (2012, September 5) retrieved 1 June 2022 from <https://medicalxpress.com/news/2012-09-clopidogrel-mi-effective-diabetes-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.