

Protection against whooping cough waned during the five years after fifth dose of DTaP

September 12 2012

OAKLAND, Calif. – Protection against whooping cough (also called pertussis) waned during the five years after the fifth dose of the combined diphtheria, tetanus, acellular pertussis (DTaP) vaccine, according to researchers from the Kaiser Permanente Vaccine Study Center. The fifth dose of DTaP is routinely given to 4- to 6-year-old children prior to starting kindergarten.

The study appears in the current online issue of the [New England Journal of Medicine](#).

This is the first study to specifically focus on the large population of highly vaccinated children who had exclusively received DTaP vaccines since birth and for whom enough time had passed since their fifth dose that DTaP vaccine waning could be measured, said the researchers. They explained that the study period included a large [pertussis](#) outbreak that occurred in California during 2010. Researchers examined the relationship between time since vaccination with the likelihood of a positive pertussis test in the Kaiser Permanente Northern California population, which includes 3.3 million members in an [integrated care](#) system with [electronic medical records](#) and a central laboratory.

Researchers compared 277 children, 4 to 12 years of age, who were positive for pertussis with 3,318 children who were negative for pertussis and separately with 6,086 matched controls. They assessed the risk of pertussis in children from 2006 to 2011 in California relative to the time since the fifth dose of DTaP and found that protection from

pertussis after the fifth dose of DTaP vaccine wanes more than 40 percent each year. The amount of protection remaining after five years depends heavily on the initial effectiveness of the fifth dose of DTaP, according to Nicola Klein, MD, PhD, co-director of the Kaiser Permanente Vaccine Study Center and the lead author of the study.

If the initial effectiveness of the fifth dose of DTaP was 95 percent, the effectiveness of DTaP would decrease to 71 percent after five years. Whereas if the initial effectiveness was 90 percent, it would decline to 42 percent after five years, explained the researchers.

"The findings suggest that whooping cough control measures may need to be reconsidered. Prevention of future outbreaks may be best achieved by developing new pertussis-containing vaccines or reformulating current vaccines to provide long-lasting immunity," said Klein.

"That said, the DTaP vaccine is effective and remains an important tool for protection against [whooping cough](#) for children and the communities in which they live, and following current CDC recommendations remains important."

The CDC currently recommends five DTaP shots for [children](#). The first three shots are given at 2, 4, and 6 months of age. The fourth shot is given at 15 through 18 months of age, and a fifth shot is given when a child enters school, at 4 through 6 years of age.

The first [pertussis vaccine](#) was developed in the 1930s and was in widespread use by the mid-1940s, when pertussis vaccine was combined with [diphtheria](#) and tetanus toxoids to make the combination whole cell pertussis vaccine DTP. In 1991, concerns about DTP safety led to the development of the acellular pertussis DTaP vaccines that are associated with fewer side effects. DTaP vaccines have completely replaced the whole cell DTP vaccines in the United States as well as in many

countries around the world.

Provided by Kaiser Permanente

Citation: Protection against whooping cough waned during the five years after fifth dose of DTaP (2012, September 12) retrieved 13 July 2023 from

<https://medicalxpress.com/news/2012-09-whooping-waned-years-dose-dtap.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.