

Mold exposure at home could increase risk for sarcoidosis

22 October 2012

People who are exposed to mold in their homes could be at an increased risk for sarcoidosis, a chronic inflammatory lung disease.

Researchers from Sweden and Slovenia tested 62 nonsmoking patients with sarcoidosis, 34 of whom had extrapulmonary manifestations (EPM).

Patients were tested for β -glucan, an immune-modulating agent found in fungi, and accompanying inflammatory biomarkers, including interleukin (IL)-6, IL-10, and IL-12.

Serum samples were also obtained from 18 [control subjects](#) with no pulmonary disease or [respiratory symptoms](#).

Results showed that levels of IL-6 and IL-12 were higher among subjects with sarcoidosis as compared with controls, and IL-12 was significantly higher among subjects with EPM. There also was a significant relation between β -glucan and mold/fungi levels in the home.

Researchers conclude that the results further support the hypothesis that exposure to fungi is important for the risk of [sarcoidosis](#).

This study was presented during CHEST 2012, the annual meeting of the [American College of Chest Physicians](#), held October 20 – 25, in Atlanta, Georgia.

Provided by American College of Chest Physicians

APA citation: Mold exposure at home could increase risk for sarcoidosis (2012, October 22) retrieved 5 May 2021 from <https://medicalxpress.com/news/2012-10-mold-exposure-home-sarcoidosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.