

# Sequential CT screening can identify indolent lung cancers

4 December 2012



Image courtesy of Blausen Medical

Changes in size on sequential low-dose computed tomography screenings, expressed as volume doubling time, indicate that about 25 percent of progressive lung cancers are slow growing or indolent, with higher lung cancer-specific mortality for new versus slow-growing or indolent cancers, according to a study published in the Dec. 4 issue of the *Annals of Internal Medicine*.

(HealthDay)—Changes in size on sequential low-dose computed tomography (LDCT) screenings, expressed as volume doubling time (VDT), indicate that about 25 percent of progressive lung cancers are slow growing or indolent, with higher lung cancer-specific mortality for new versus slow-growing or indolent cancers, according to a study published in the Dec. 4 issue of the *Annals of Internal Medicine*.

Giulia Veronesi, M.D., from the European Institute of Oncology in Milan, and colleagues assessed VDT as an indicator of overdiagnosis for screening-detected lung cancer by estimating the VDT in 175 high-risk patients who were diagnosed with primary lung cancer. VDT was characterized as fast growing (VDT

APA citation: Sequential CT screening can identify indolent lung cancers (2012, December 4) retrieved 3 October 2022 from <https://medicalxpress.com/news/2012-12-sequential-ct-screening-indolent-lung.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no*

*part may be reproduced without the written permission. The content is provided for information purposes only.*