

## Heavy moms-to-be at greater risk of Csection

5 March 2013

Researchers from Norway found that women with a pre-pregnancy body mass index (BMI) of 40 had an increased risk of vacuum extraction delivery or Cesarean section (C-section). Findings that appear in *Acta Obstetricia et Gynecologica Scandinavica*, a journal published by Wiley on behalf of the Nordic Federation of Societies of Obstetrics and Gynecology, indicate that women with more than a 16 kg (30 lbs) weight gain during pregnancy increased their risk of forceps or vacuum extraction, and C-section.

Obesity is a global health crisis, with the <u>World</u> <u>Health Organization</u> (WHO) reporting that 1.4 billion adults were overweight (BMI of 25-29.9) in 2008 and more than half a billion would be considered obese (BMI of 30 or more). Roughly one-third of U.S. adults were considered obese in 2009-2010, according to the <u>Centers for Disease</u> <u>Control and Prevention</u> (CDC). In Norway previous studies estimate that <u>obesity prevalence</u> among women of reproductive age has increased two- to three-fold in the last generation.

"Our study examines pre-pregnancy BMI and gestational <u>weight gain</u> on the mothers' risk of operative delivery," said Dr. Nils-Halvdan Morken from the University of Bergen in Norway and lead author of the present study. "With such alarming rates of obesity understanding its impact is an important health issue, particularly for women in child-bearing years."

The research team used data taken from participants of the Norwegian Mother and Child Cohort Study (MoBa)—a prospective, populationbased group of women giving birth in Norway between 1999 and 2008. The MoBa study, conducted by the Norwegian Institute of Public Health, now includes a cohort of 108,000 children, 90,700 mothers, and 71,500 fathers. A sample of 50,416 women who gave birth to one child were included in the study and those experiencing preeclampsia, hypertension, diabetes, <u>gestational</u> diabetes or placenta previa were excluded.

Results show that overweight and obese women before pregnancy were at increased risk of Csection. Women with a pre-pregnancy BMI of 40 or more had the strongest risk of C-section and increased risk of vacuum extraction delivery. Researchers also found that women who gained 16 kg or more while pregnant significantly increased the risk of forceps, vacuum extraction and Csection—a finding that was independent of BMI prior to pregnancy. Obese women had significantly lower gestational weight gain, but their babies tended to be larger.

Dr. Morken concludes, "Obesity and weight gain above 16 kg during pregnancy are independent risk factors for vacuum extraction delivery and need for C-section. While other factors may contribute to operative delivery and further investigation of gestational weight gain is warranted, it is important obstetricians be aware of the impact of a high BMI on pregnancy and delivery to properly advise women considering motherhood."

**More information:** "Pre-Pregnant Body Mass Index, Gestational Weight Gain and the Risk of Operative Delivery." Nils-Halvdan Morken, Kari Klungsoyr, Per Magnus, Rolv Skjarven. *Acta Obstetricia et Gynecologica Scandinavica*; Published online: February 19, 2013 (DOI: <u>10.1111/aogs.12115</u>)

Provided by Wiley



APA citation: Heavy moms-to-be at greater risk of C-section (2013, March 5) retrieved 11 October 2022 from <u>https://medicalxpress.com/news/2013-03-heavy-moms-to-be-greater-c-section.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.