

Sex between monogamous heterosexuals rarely source of hepatitis C infection

19 March 2013

Individuals infected by the hepatitis C virus (HCV) have nothing to fear from sex in a monogamous, heterosexual relationship. Transmission of HCV from an infected partner during sex is rare according to new research published in the March issue of *Hepatology*, a journal published by Wiley on behalf of the American Association for the Study of Liver Diseases (AASLD).

Experts estimate that HCV affects up to 4 million Americans, most of whom are sexually active. Medical evidence shows HCV is primarily transmitted by exposure to infectious blood, typically through intravenous (IV) drug use. However, there are conflicting reports regarding sexual activity and HCV transmission with some studies suggesting that exposure to infected blood during sex—through bodily fluids such as vaginal secretions, semen or saliva—may carry a minimal infection risk.

"Generally the risk for transmitting HCV to sex partners is very low," explains lead study author Dr. Norah Terrault with the University of California, San Francisco. "Yet, lack of [quantitative data](#) about the risk of HCV transmission with sexual activity remains a limitation for doctors counseling their patients on safe [sex practices](#)."

To specifically quantify the risk HCV transmission from a chronically infected individual to their sex partner, researchers recruited 500 anti-HCV-positive individuals, who were negative for the human immunodeficiency (HIV), and their long-term heterosexual partners. Couples were surveyed about [lifetime risk](#) factors for HCV infection, [sexual practices](#) of the couple, and sharing of personal items. The team analyzed blood samples to determine the presence or absence of active virus in the blood and compared the HCV strains in those couples with HCV present.

The majority of HCV infected individuals who

participated in the study were non-Hispanic whites, had a median age of 49 years, and sexual activity with their partners ranging from 2 to 52 years. HCV prevalence among partners was 4%, with 9 couples having similar viral strains and viral samples from 3 couples were highly related which is consistent with HCV transmission between the partners.

The maximum incidence rate of HCV transmission by sex was 0.07% per year or roughly 1 per 190,000 sexual contacts that researchers based upon 8377 person-years of follow-up. The team did not identify any specific sexual practices linked to HCV infections among the couples. "Our study provides clinicians with important information for counseling chronic HCV patients in long-term sexual relationships, supporting the current recommendations that couples not change their sexual practices if they are in a monogamous heterosexual relationship," concludes Dr. Terrault.

More information: "Sexual Transmission of HCV Among Monogamous Heterosexual Couples: The HCV Partners Study." Norah A. Terrault, Jennifer L. Dodge, Edward L. Murphy, John E. Tavis, Alexi Kiss, T.R. Levin, Robert Gish, Michael Busch, Arthur L. Reingold, Miriam J. Alter. *Hepatology*; (DOI: [10.1002/hep.26164](https://doi.org/10.1002/hep.26164)); Print Issue Date: March, 2013.

Provided by Wiley

APA citation: Sex between monogamous heterosexuals rarely source of hepatitis C infection (2013, March 19) retrieved 27 April 2021 from <https://medicalxpress.com/news/2013-03-sex-monogamous-heterosexuals-rarely-source.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.