

# Antidepressants for pregnant moms don't affect infants' growth, research says

20 March 2013

Selective serotonin reuptake inhibitor (SSRI) antidepressants taken by a woman during pregnancy do not impact her infant's growth over the first year, reports a new study from a Northwestern Medicine scientist.

There had been concern that antidepressant treatment during pregnancy reduced growth during the first year. Previous data suggested depression during pregnancy also could diminish infant growth.

But the new study showed infants born to mothers who took SSRIs during pregnancy had a similar weight, length and [head circumference](#) over the first year as infants born to non-depressed women who did not take antidepressants. The infants whose mothers took antidepressants were shorter at birth, but the difference disappeared by two weeks of age.

In addition, growth measurements for the infants of [depressed women](#) who did not take SSRIs were similar to the general population.

"Most women want to know about the effect of their depressive illness or the medication they take during pregnancy not only on the infant at birth, but also on the baby's longer-term growth and development," said Northwestern Medicine lead author Katherine L. Wisner, M.D. "This information may help women balance the risks and benefits of continuing their antidepressant treatment during pregnancy."

Wisner is director of Northwestern's Asher Center for the Study and Treatment of Depressive Disorders. She also is the Norman and Helen Asher Professor of Psychiatry and Behavioral Sciences and of [obstetrics and gynecology](#) at Northwestern University Feinberg School of Medicine and a physician at Northwestern Memorial Hospital.

The study will be published March 20, 2013 in *The American Journal of Psychiatry* in Advance, its online ahead-of-print website.

Depression has negative impact on a mother's and infant's health, Wisner said, noting that women who stop taking SSRIs near the time of conception have a high relapse rate.

Maternal prenatal stress and depression are linked to preterm birth and low infant birth weight, which increases the risk of cardiovascular disease. Depression also affects a woman's appetite, nutrition and prenatal care and is associated with increased alcohol and drug use. Women with untreated depression have a higher body mass index preconception, which carries additional risks.

Provided by Northwestern University

APA citation: Antidepressants for pregnant moms don't affect infants' growth, research says (2013, March 20) retrieved 3 May 2021 from <https://medicalxpress.com/news/2013-03-antidepressants-pregnant-moms-dont-affect.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*