

# Hospital infections in cancer patients climbed, deaths declined, study finds

26 March 2013

In a nationwide study of patients undergoing surgery for cancer, Henry Ford Hospital researchers have found that while infections during hospital stays increased during a 10-year period, the death rate from those infections dropped.

The findings suggest that diagnosis and management of healthcare-associated infection, or HAI, have improved over time.

The study appears in the current issue of the journal *Cancer*.

"[Cancer patients](#) often have surgery as part of their treatment and are at increased risk for developing healthcare-associated infection," says study lead author Jesse Sammon, D.O., with Henry Ford's Vattikuti Urology Institute.

"Ironically, they have often been left out of previous studies of healthcare-associated infection following surgery precisely because they're at higher risk of developing these infections."

HAIs are one of the most common and dangerous adverse consequences of hospitalization – but they are preventable. Under pressure from growing public awareness, at least 27 states have enacted laws as of 2010, requiring [health systems](#) and hospitals to report HAI rates.

This increased public awareness was due in large part to the 1999 publication of the Institute of Medicine's landmark report; *To Err is Human*, which compelled medicine to examine the depth and breadth of the HAI problem. And the [medical field](#) responded by adopting numerous guidelines and programs to better detect and prevent such infections.

The Henry Ford study suggests that those initiatives are working.

Drawing on the Nationwide Inpatient Sample

database, researchers identified more than 2.5 million [patients](#) who had major [cancer surgery](#) for one of eight types of malignancies between 1999 and 2009. These included colectomy, cystectomy, esophagectomy, gastrectomy, [hysterectomy](#), pneumonectomy, pancreatectomy and [prostatectomy](#).

Among the patients, Dr. Sammon and his colleagues found that the incidence of HAI related to these surgeries increased 2.7 percent per year during the 10-year study period; but at the same time, the rate of deaths linked to the HAIs dropped 1.3 percent per year.

While those results are encouraging, Dr. Sammon said they are tempered by additional findings that socio-demographic factors such as race and insurance status continue to be serious national health policy concerns.

When compared to white patients, African Americans showed a 26 percent increase in HAI.

And if a patient's health care payments were made through Medicare or Medicaid – or if the patient was uninsured – the odds of HAI went up 18 percent to 67 percent.

Provided by Henry Ford Health System

APA citation: Hospital infections in cancer patients climbed, deaths declined, study finds (2013, March 26) retrieved 15 November 2022 from <https://medicalxpress.com/news/2013-03-hospital-infections-cancer-patients-climbed.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*