


# Poor 'Health literacy' keeps patients from taking meds

27 March 2013



In study, adults with diabetes who didn't grasp health info often skipped needed drugs.

(HealthDay)—Adult diabetes patients who don't understand basic health information are less likely to continue taking newly prescribed antidepressants, a new study finds.

This is an important issue because depression in adults with diabetes is often chronic and may require long-term treatment with medication, the researchers said.

The nearly 1,400 patients in the study were followed for 12 months after being prescribed an antidepressant. Most of the patients filled the prescription at least once, but 43 percent did not refill the prescription and nearly two-thirds had stopped taking their [antidepressant medication](#) by the end of the study.

The investigators found that 72 percent of the patients struggled to understand basic health instructions, which the study authors called "limited [health literacy](#)." These patients were much less likely to keep taking their [antidepressants](#) than those with [good health](#) literacy, the authors said in a news release from Kaiser Permanente.

This difference was not explained by other factors

known to be associated with patients not taking prescribed medications, including age, race and ethnicity, income, and level of English-language skills, according to the study, which was published in the March issue of the *Journal of General Internal Medicine*.

"The high rates of early discontinuation ... among adults with diabetes who had any health literacy limitation suggest that few of these individuals received an adequate course of antidepressant therapy," lead author Dr. Amy Bauer, of the University of Washington School of Medicine, said in the news release. "Getting that sufficient treatment is critical in preventing [relapse](#) and recurrence of depression."

"Physicians should be aware of this," Bauer added. "For antidepressant treatment to succeed, patients with limited health literacy may require more intensive counseling and clearer explanations about use of antidepressant medications and closer follow-up."

Study senior author Andrew Karter, a research scientist at Kaiser Permanente, said other studies have found that "those with health literacy limitations are more likely to have poor control of their chronic medical conditions, such as diabetes, congestive heart failure and HIV."

He added, however, that "this is the first study to examine the association between health literacy and antidepressant adherence among patients with diabetes. This type of research gives our health care systems important feedback because, as providers, we often remain unsure whether the critical health information we convey to our patients is fully understood."

The researchers said the new findings highlight the importance of efforts to improve people's health literacy, simplify information about treatment options, improve public understanding about

depression treatment, and monitor whether [patients](#) are taking their prescribed antidepressants.

**More information:** The U.S. National Institute of Mental Health has more about [diabetes and depression](#).

[Health News](#) Copyright © 2013 [HealthDay](#). All rights reserved.

APA citation: Poor 'Health literacy' keeps patients from taking meds (2013, March 27) retrieved 6 May 2021 from <https://medicalxpress.com/news/2013-03-poor-health-literacy-patients-meds.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*