

# Risk of kidney disease doubled with use of fluoroquinolone antibiotics

3 June 2013

The risk of acute kidney disease is doubled for people taking oral fluoroquinolone antibiotics, according to a study published in *CMAJ* (*Canadian Medical Association Journal*).

Fluoroquinolones, including ciprofloxacin, levofloxacin and moxifloxacin, are common broad-spectrum antibiotics most often used to treat respiratory and urogenital infections. Case reports have indicated acute kidney injury with use, and prescription labels carry a warning of [kidney failure](#). However, when oral fluoroquinolones are prescribed in clinical practice, kidney injury is usually not considered.

Researchers from the US and Canada undertook a study to determine the risk of acute kidney injury with the use of oral fluoroquinolones for adult men 40

**More information:**

[www.cmaj.ca/lookup/doi/10.1503/cmaj.121730](http://www.cmaj.ca/lookup/doi/10.1503/cmaj.121730)

Provided by Canadian Medical Association Journal

APA citation: Risk of kidney disease doubled with use of fluoroquinolone antibiotics (2013, June 3) retrieved 30 July 2022 from <https://medicalxpress.com/news/2013-06-kidney-disease-fluoroquinolone-antibiotics.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*