


Lixisenatide non-inferior to exenatide for diabetes

7 June 2013



In adults with type 2 diabetes not adequately controlled with metformin, lixisenatide once daily offers diabetes control similar to that of exenatide twice daily, with some added benefits, according to a study published online May 22 in *Diabetes Care*.

(HealthDay)—In adults with type 2 diabetes not adequately controlled with metformin, lixisenatide once daily offers diabetes control similar to that of exenatide twice daily, with some added benefits, according to a study published online May 22 in *Diabetes Care*.

Julio Rosenstock, M.D., from the Dallas Diabetes and Endocrine Center at Medical City, and colleagues randomly assigned 634 adults whose type 2 diabetes was not adequately controlled with metformin ([hemoglobin A1c](#) [HbA1c], 7 to 10 percent) to 20 µg lixisenatide once daily or 20 µg exenatide twice daily.

After 24 weeks, the researchers found that lixisenatide once daily was non-inferior to exenatide twice daily in terms of least squares mean change in HbA1c, responder rate (HbA1c

APA citation: Lixisenatide non-inferior to exenatide for diabetes (2013, June 7) retrieved 15 June 2021 from <https://medicalxpress.com/news/2013-06-lixisenatide-non-inferior-exenatide-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no

part may be reproduced without the written permission. The content is provided for information purposes only.