

# Big belly increases death risk in heart attack survivors

1 September 2013

Having a big belly increases the risk of death in heart attack survivors, according to research presented at ESC Congress 2013 by Professor Tabassome Simon and Professor Nicolas Danchin from France. The findings from the FAST-MI 2005 registry suggest that lifestyle interventions in heart attack patients should focus on losing abdominal fat.

Professor Simon said: "The impact of obesity on long term mortality and [cardiovascular complications](#) in the general population has been the object of recent debate. Much emphasis has also been given to the deleterious role of abdominal obesity."

She added: "At the time of a heart attack, early mortality tends to be lower in [obese patients](#), a phenomenon well known in critical care situations and described as the 'obesity paradox'. Little is known, however, about the potential impact of obesity and abdominal obesity on long-term outcomes in patients who have survived the acute stage of a heart attack."

The current study sought to determine the associations between [body mass index](#) (BMI, kg/m<sup>2</sup>) and [waist circumference](#) with 5-year mortality in patients of the FAST-MI 2005 registry who had left the hospital alive.

FAST-MI 2005 is a French nationwide survey which included 3,670 patients from 223 institutions who were admitted for [acute myocardial infarction](#) to an [intensive care unit](#) at the end of 2005. Long-term follow-up was achieved in 99.6% at one year, 98% at 3 years, and 95% at 5 years. Of the 3,463 patients who were discharged alive, BMI was recorded in 3,102 and waist circumference in 1,647 patients. Statistical techniques were used to take into account the differences in baseline characteristics between the different BMI groups.

At 5 years, absolute mortality was highest in the

leanest patients (BMI 100 cm in women and >115 cm in men) was also associated with increased long-term mortality.

Professor Simon said: "As waist circumference is strongly linked to BMI, we determined the upper quartile of waist circumference (i.e. quarter of the population with the highest waist circumference) within each BMI category and used both variables together to determine their respective role in association with long-term mortality."

She continued: "We found that both lean patients (BMI

APA citation: Big belly increases death risk in heart attack survivors (2013, September 1) retrieved 2 May 2021 from <https://medicalxpress.com/news/2013-09-big-belly-death-heart-survivors.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*