

Bacteria responsible for gum disease facilitates rheumatoid arthritis

12 September 2013

Does gum disease indicate future joint problems? Although researchers and clinicians have long known about an association between two prevalent Studies indicate that compared to the general chronic inflammatory diseases - periodontal disease and rheumatoid arthritis (RA) - the microbiological mechanisms have remained unclear.

In an article published today in PLOS Pathogens, University of Louisville School of Dentistry Oral Health and Systemic Diseases group researcher Jan Potempa, PhD, DSc, and an international team of scientists from the European Union's Gums and Joints project have uncovered how the bacterium responsible for periodontal disease, Porphyromonas gingivalis worsens RA by leading to earlier onset, faster progression and greater severity of the disease, including increased bone and cartilage destruction.

The scientists found that P. gingivalis produces a unique enzyme, peptidylarginine deiminanse (PAD) which then enhances collagen-induced arthritis (CIA), a form of arthritis similar to RA produced in the lab. PAD changes residues of certain proteins into citrulline, and the body recognizes citullinated proteins as intruders, leading to an immune attack. In RA patients, the subsequent result is chronic inflammation responsible for bone and cartilage destruction within the joints.

Potempa and his team studied another oral bacterium, Prevotella intermedia for the same affect, but learned it did not produce PAD, and did not affect CIA.

"Taken together, our results suggest that bacterial PAD may constitute the mechanistic link between P. gingivalis periodontal infection and rheumatoid arthritis, but this ground-breaking conclusion will need to be verified with further research," he said.

Potempa said he is hopeful these findings will shed

new light on the treatment and prevention of RA.

population, people with periodontal disease have an increased prevalence of RA and, periodontal disease is at least two times more prevalent in RA patients. Other research has shown that a P. gingivalis infection in the mouth will precede RA, and the bacterium is the likely culprit for onset and continuation of the autoimmune inflammatory responses that occur in the disease.

Provided by University of Louisville



APA citation: Bacteria responsible for gum disease facilitates rheumatoid arthritis (2013, September 12) retrieved 30 June 2022 from https://medicalxpress.com/news/2013-09-bacteria-responsible-gum-disease-rheumatoid.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.