

Longer detox might work better for prescription pain med addiction

23 October 2013



Small study compared varying treatment periods used with opioid-dependent patients.

(HealthDay)—A longer period of detoxification may be more effective for people being treated for addiction to prescription painkillers called opioids, according to a small new study.

Abuse of prescription opioids such as oxycodone, hydrocodone and hydromorphone is a major public health problem in the United States. The new 12-week study, which included 70 people undergoing outpatient treatment for opioid addiction, was published online Oct. 23 in the journal *JAMA Psychiatry*.

For the first two weeks, all the patients took buprenorphine, a medication used to treat opioid addiction. They were then randomly assigned to slowly reduce the dose of buprenorphine over one, two or four weeks, followed by treatment with naltrexone, a medication that blocks opioid strength.

Patients in the four-week group were more likely to stop abusing opioids than those in the one-week or two-week groups, according to a journal news release.

The findings suggest that some prescription opioid abusers may respond positively to outpatient treatment with [buprenorphine detoxification](#) followed by naltrexone while undergoing behavioral

therapy [counseling], study authors wrote.

"Additional controlled studies are needed to better understand the parameters of efficacious treatments for [prescription [opioid](#)] dependence, as well as to identify the individuals for whom brief vs. longer-term treatments are warranted," concluded study authors Stacey Sigmon, of the University of Vermont, in Burlington, and colleagues.

More information: The U.S. National Institute on Drug Abuse has more about [prescription drug abuse](#).

Copyright © 2013 [HealthDay](#). All rights reserved.

APA citation: Longer detox might work better for prescription pain med addiction (2013, October 23) retrieved 9 July 2022 from <https://medicalxpress.com/news/2013-10-longer-detox-prescription-pain-med.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.